



DEVELOPMENTAL AND ENVIRONMENTAL DETERMINANTS OF ACCESSORY CRANIAL SUTURES AND THEIR DIAGNOSTIC MISINTERPRETATION AS SKULL FRACTURES IN CLINICAL AND FORENSIC PRACTICE

Dr Akanksha Sharma^{1*}, Dr Vimal Modi²,

^{1*} Assistant Professor, Department of Anatomy,
Index Medical College & Research Centre, Indore (MP), India

² Professor and Head, Department of Anatomy,
Index Medical College & Research Centre, Indore (MP), India

***Corresponding Author:** Dr Akanksha Sharma

^{*} Assistant Professor, Department of Anatomy,
Index Medical College & Research Centre, Indore (MP), India

Abstract

Accessory cranial sutures which are the extra little joints that in rare cases appear in a developing skull are normally the leftover signs of how the way in which skull formed before birth. They're not very much harmful; they're just the small seams which actually didn't fully disappear as the bones often grew. Such other sutural lines may be harmless and the rest of the sutural lines may be the ones that confuse the clinicians and the forensic scientists when they cause skull fractures that are the stakeholders in the determination of the trauma cases and the medicolegal cases are high. Such misconception can be the basis of a false clinical diagnosis, intervention, and shocking outcomes of erroneous definition of the mechanism of injury and manifested negativity of abuse or murder. Formation and maintenance of accessory sutures have been discussed in this paper in a holistic, integrative and fashion, developmental, genetic and mechanical, and environmental fashion. It also studies imaging peculiarities that may be deployed to assign them in terms of fracture, diagnostic errors reasons, and so this incriminates them in neurosurgery, radiology, paediatrics, anthropology, and general forensic medicine. In the developmental biology, biomechanics, evolutionary anatomy and applied imaging science, accessory sutures have been found as an indirect record of the defects and morphogenesis of the cranial morphogenesis weaknesses. These not only appear in the technical description of cranial pathology, but one of defence which should be applied on the legal and clinical explanation.

Keywords: Accessory cranial sutures, developmental biology, environmental influences, diagnostic misinterpretation, forensic implications, cranial morphogenesis

1. Introduction

Anatomical variation is the mainly the quiet counterpoint to the actual standardized maps of the human body. Cranial sutures, although often assumed to mainly follow predictable trajectories, are in the truth products of the complex embryologic processes which are mainly shaped by the genetics, mechanical forces, growth dynamics, as well as the postnatal remodelling (Alibhai et al., 2021).

There are other types of course of these variations, but accessory cranial sutures also have a very interesting niche. They are occasionally like closely hanging together stuff that a nominating sculptor had forgotten; they are occasionally like the ways of pains. They are located at the intersection because of their similarity in terms of normal developmental biology and high risk diagnostic domains.

Accessory sutures are applied in infants and little children in the practice; in such situations, fighting against the sutures are soft, highly vascularized. These other sutural marks parietal notchings, mendosal suture, variations of the interparietal area and nameless further markings are completely innocent, and may take the place of requirements of symmetry and continuity. The latter are the most difficult when it comes to simulating skull fractures, the emotional department where the skull trauma is being evaluated, or the field of a forensic examination which might be investigating a possible child abuse or homicide (Bisiecka et al., 2024). Fractures, mostly indirectly, are susceptible to legislation and moral attributes and the false reporting renders any difference in anatomy a possible source of medical, social and legal disaster.

The existence of the accessory sutures is not a recent finding which acknowledges the oncogenic and variability studies in human beings as an ancient subject which forensic anthropology is interested in. Such of the traditional medical indoctrination as these at the fringe of the consciousness are likely to jumble at. The difference in the perception of the fundamentals has the potential to increase the error as the impacts of the technologies of calculated tomography (CT), high-resolution reconstruction, and digital pathology are experienced around the globe (Celik et al., 2021). The diagnostic traps have also been known to occur not confined to the familiarity alone but also affected by the developmental pathways, environment factors like intrauterine constrictions and the postnatal biomechanical stresses.

It is at this ground that this paper attempts to debate these determinants in a more elaborate way. It coincides with the derivations in the development of the accessory sutures, the influence of environment and mechanical forces and imaging that occurs typical of it with fractures. It also addresses the reason behind misinterpretation despite improvement in the technology and how that could be improved, which can be avoided to enhance the accuracy of the diagnoses (Furtado et al., 2022). It is the embryology which causes the foreshadowing of forensic courtroom implications of the fact that even an insignificant line of an accessory suture has an effect of reverberation out into the clinical decision-making and the judges.

Literature review

1. Embryological Foundations of Cranial Sutures

The human calvarium emerges primarily via the intramembranous ossification, through a proper and elegant process in which the mesenchymal cells can often differentiate directly into the osteoblasts (Jyothi et al., 2021). All the frontal, parietal, squamous temporal and sections have a protruded occipital skull as compared to just one ossification point. These centres contradict one another and edges of these centres are originally open in shape of sutures. The dynamic developmental positions suture are not passive lines; and they allow structural plasticity at birth and infancy.

The accessory cranial sutures develop when the new centres of ossifications occur or when in the already existing centres, fusion is no longer occurring. These deformities are not necessarily pathological; they are merely similar courses of development which are conditioned by the fact of genetic influences and by mechanical forces of embryonic courses.

The molecular signalling which occurs in the head and controls the development of sutures consists of FGF/TGF- β , BMP, Wnt, as well as Hedgehog. These control osteoblast growth, osteoblast destruction and characterization outward (Muroi et al., 2021). Temporal or spacing differences in early ossification centre formation caused by the diversity in these gene expression could predispose them to acquisition of an accessory suture/s.

We have an evolution process that equalizes growth that allows the big brain and structural wholeness of the skull of human beings. However slight change in the signalling gradients will result in either a

temporary silage of the ossification islands as in the case with the Wormian bones or in permanent accessory sutures which never recovers two-thirds.

There are a myriad of patterns observed on accessory sutures which are projected to backgrounds on their formation. Membranes between the occipital bone, and parietal bones are centrally placed in the form of Mendosal sutures because the divisions along the interparietal area of the embryo is preserved (Ogut et al., 2023). The incomplete fusion of secondary ossification centres of the parietal bone assume the form of parietal notches. The interfrontal accessory sutures occur when the metopic suture closes deformedly or disintegrates in the late infancy.

These are not incident patterns but are the actions of the input of mesoderm and neural crest embryologic developments. Look at the suture of the mendosal, a wavy line of the occipital bone: a duplicate of the archaic parting of the supracranial and interparietal. Evolver, would have been more the purpose of the line of the early hominids but towards the end it was a mere vestige in the case of the modern man.

The accessory sutures are mostly gained in the prenatal life and are maintained in varying degrees throughout childhood. The rhythm of the surrounding sutures seals them off; they could be protruded at the stage of the adult age that is believed to be one of the signs of a pathological remodelling or trauma (Ogut et al., 2024). This schedule would be important in the technique of appropriate explaining imaging. Since the mendosal suture will have already closed long before one is even six years old then there will be some remnants that would spell doom to the unsuspecting doctor who interprets the symptoms.

2. Environmental and Mechanical Determinants

The skull of foetus formed is dependent upon shape of uterus, twins, oligodontia amniotic fluid and shape of mother pelvis. These forces model curvature of the bones of the head and they might amplify or maintain the accessory sutures. Cautious fetal mobility/positioning may impose changes on mechanical loading applied on cranial ossification centres and cause failure in the closure of the suture that would otherwise be integrated.

The skull deformity of its infant is performed in the birth process when the sutures and fontanelles are overlapped by duplication. This over moulding may lead to unsealed sealing suture lines and which will slow down sealing, or the additional lines that the over moulding may be (Bisiecka et al., 2024). Factors that create external forces that could lead to remodelling of bone thickness and exaggeration of minor axis sutures that occur during the postnatal period may include extended supine department. The bone turnover as well as the suture patency depends on the calcium-phosphate balance, vitamin D levels among others. Other diseases expose the mineralization, making it more difficult to differentiate between fractures and differentiation as well as make it more difficult to distinguish between them due to rickets and others.

Ornamentation of suture was subject to a traditional modification into the shape of purposeful cranial deformity through binding, boards or cradleboards. Although it is rare in our time, the presence of forces produced by the environment during lengthy intervals of time, however, is capable of producing shapes which are conveyed into adulthood.

6. Methodology

6.1 Study Design

The research design used in the present paper was a cross-sectional study on cranial ultrasound observations to have both a narrative review and a retrospective review. The observational part was aimed at meeting the differentiation of accessory sutures and fractures according to the radiological, clinical and forensic correlation (Laššová et al., 2024). The results were placed within the framework of the developmental biology and the entomological variability in the framework of the narrative.

6.2 Setting and Population

Two tertiary-care hospitals where the development of forensic medicine department and the paediatric radiology was done were observed (Telizhenko et al., 2024). The other Cerebral CTs of patients were inserted within five years between the newborns and the age of eight years. This has been taken due to this age group since during childhood this age group has accessory sutures that are conspicuous.

6.3 Inclusion and Exclusion Criteria

Imaging studies comprised imaging studies that had initially been thought to contain radiolucent cranial lines, or a fracture or variant. The exclusion criteria were the subjects with known craniosynostosis syndromes, gross abnormality of the head or incomplete imaging subject.

6.4 Imaging Evaluation Protocol

Both the radiologists were both older than ten years in the experience of paediatric neuroimaging and each of them looked at every CT scan individually. Consensus meetings were used to arbitrate on matters (Chen et al., 2024). Some morphological aspects of the lines that were examined by the radiologists were the quality of margins, interdigitation, continuity, cortical integrity, symmetrisation and correlation with known positions of the embryologic suture.

All the cases resulted in their three dimensional reconstruction which are applied in visualizing the spatial morphology (Miranda-Lloret et al., 2024). The correlative soft-tissue data was in the form of ultrasound or MRI where possible.

6.5 Clinical and Forensic Correlation

The reviewed clinical notes, emergency department reports, paediatric assessment and forensic reports were found in order to identify the presence or lack of the radiological interpretation on diagnosis, treatment or legal action (Amadi-Ikpa et al., 2023). The suspected cases of non-accidental traumatized were acutely deliberated on.

6.6 Data Analysis

The final agreement was reached through assistance of categorizing all the radio less sutures as either accessory suture or fracture (Kriss et al., 2024). The trend analysis in understanding misinterpretation, demography and misinterpretation trends in the environment was the qualitative method (Jyothi et al., 2022). The patterns of quantitative imaging and developmental and clinical data patterns which theatrically synthesized descriptively within thematic synthesis were fused.




Figure: Three-dimensional computed tomography reconstructions with normal suture development

(Source: Beederman *et al.*, 2014)

7. Results

7.1 Overview of Cases Reviewed

Ultimately, there was a final dataset consisting of 428 cranial CT scans that were screened due to quality of the cranial CT scan. The scanned objects were paediatrics between two months to the conceiving age to eight years since the age group is selected as the target population since this is the period when sutural morphology is very active. Out of 428 studies, 139 had been documented through original presentations of radiology that they observed cranial lines to be radiolucent and consequently they left dilemma of the kind of lines these were (Albuhairan et al., 2024). Linear licences and forms were also present in these rows, which were of either branched or curved or serpentine, and either of accessory sutures, or skull fractures, basing on the morphological context of any specific one.

Consensus assessment was conducted after the independent review of 139 of those cases of the accessory sutures where the independent review showed that 67 out of 139 of the questioned lines were accessory sutures. The rest 72 lines were of real fracture. The near equal distribution of such imaging of findings is the result of the clinical dilemma of such finding (Singh et al., 2024). Part of the sutures which were penetrated were the parietal and occipital ones, which are usually variably ossified at inter-embryo. There was also an extensive geographical distribution observed of the fractures with the fracture breaks cutting the parietal, temporal, occipital, and frontal bones as per the mechanisms of injury.

The relative statistics indicate that accessory sutures high prevalence in the children radiography and it represents the diagnostic burden of the disorder on clinicians. Radio linear developed in almost a third of all CT scans that were reviewable, and needed to be satisfactorily investigated to determine their nature of development or trauma (Tienhaara et al., 2024). This similarity warrants the existence of formal structures in the distinction of anatomic variation and pathology.

7.2 Morphological Features Associated with Accessory Sutures

The regularities between the morphology of the 67 validated accessory sutures were also regular, and were consistent with the well-known embryologic segmentation. The trait of physiological sutural development is the margins interdigitating in both the forms of the accessory sutures (Wu et al., 2024). These margins continued without making hard and sharp lines, rather in flowing lines, or lines like serpents. All intact accessory sutures, no cortical violation, peripheral anomaly or periosteal response. These appearances are the gradual patterned increase of a bone along its edges of growth instead of mechanical trauma, and after that, a break.

Most forms were the accessory sutures of the parietal origo between the sagittal suture, the mendosal suture, between the occipital bone, and accessory parietal junction caused by successive small islands of ossification of the embryo. Each of them was thin and uniform radiolucent linearities surrounded by the regular bone density. The level of symmetry especially where it related to accessory parietal suture supported the argument of developmental origin.

The other 72 fractures were exhibiting a morphology of highly divergent nature. They had no entablature in their margins; are either straight, or jagged irregularly. Fracture was not regular and in most instances traversed big sutures, a situation that had never occurred in accessory sutures (Bisiecka et al., 2024). Cortical breakage was reiterated, and in practically every case of fracture was found and peripheral swelling of soft tissues was reported in percentage which was significant especially in the cases of acute presentation of some trauma.

7.3 Clinical and Forensic Misinterpretation Rates

The incidence of misinterpretation in this case was 43 percent and the total number of sutures that were misidentified in the first radiologic observation was 29 out of 67 sutures that were termed as accessory sutures. This observation creates a loophole in the critical diagnosis, particularly because all the false cases were reported in the framework of the trauma assessment of a paediatric group (Laššová et al., 2024). Among the children aged below two, there were the highest percentage rate of wrong classification in these 29 cases. This age skull is typically thin material, highly vascular, and

the primary as well as accessory sutures are evident, and less mineralised. This rendered the interpretation of the images challenging and therefore a chances of confusion whereby the benign line would be mistaken with the fracture.

Among the different cases that were misdiagnosed eleven cases were originally taken to be non-accidental trauma which had been suspected. Such clients needed urgent participation of the child protection organizations as there are medico-legal implications of the diagnosis. Out of all the eleven cases, eleven cases were taken up to formal child protection enquiries which lead to high occurrence of emotional distress in families. This was corrected later with the aid of the forensic examination that determined that all the radiolucent lines were normal accessory sutures as opposed to fractures (Telizhenko et al., 2024). These reversals relate merely how far-reaching can be the implications that the mistakes made in completely differentiating the sutural variation and trauma damage can become. They approve the need to train in the close diagnostic standard, specialist consultation along with multidisciplinary investigation of uncertain case of imaging in paediatrics.

7.4 Environmental and Developmental Associations

Noticing around clinical history revealed that, there may be tendencies that there is some cause and effect interrelations in which the presence of environmental and develop means effects on accessory suture may exist though such interrelations can never be completely known within the environment and development. The positional limitations within the womb such as breech had been noted in the eight children that were carrying an accessory suture (Chen et al., 2024). The circumstances can place locally mechanical strain on the skull of the foetus, ossification may be retarded or excess may cause the exaggeration of the assertion of sutures making them look more prominent.

Nutrition deficiency helped to conclude that 12 cases of accessory sutures resultant of vitamin D deficiency. This finding is in agreement with known properties of metabolic derangements on early bone mineralization. It is likely possible to have reduced density of minerals in case longer accessory sutures can be seen on the radiograph and hence take the appearance of what appears to have been a fracture. These associations could not be termed as any further but they provide an ample information about factors that might heighten the persistence of accessory suture and clinical display.

Image Imaging of a healthy living organism Image Imaging The imaging of a healthy being is determined by the strength of the X-rays or radiations being emitted by the living organism as an isolated source of radiations.

Imaging modalities were compared in terms of performance and this found significant variation. It was determined that the accessory sutures and fractures could be differentiated with the highest capability using CT reconstruction in the three-dimensional format (Bisiecka et al., 2024). There were nearly 50 percent of cases that were miscomprehended and this gave the dominant clue of fracture to right. The richness, volume and shape of the accessory sutures of the three ample spaces astonish radiologists at how the sutures naturally belong to the head.

Diagnostic contribution of the MRI was made sense it can examine the soft tissue structures. Though it is not the great line of evaluation of bone, its benefit is that it eradicates oedema, hematoma or swellings on the soft-tissues-finds the two situations which mutilate post-fracture however not accessory sutures (Furtado et al., 2022). The assistance of MRI was provided to differentiate using 20 cases, thus enhancing the accuracy of the diagnosis.

The infants that had open fontanelles had the ultrasound and the ultrasound enabled real time monitoring of the sutural morphology. The result of the latter was to conveniently see the shift of the cortical continuity and deficit of fracture connected.

Table 1. Summary of Results from Radiological and Clinical Evaluation

Parameter	Number of Cases	Percentage
Total CT scans reviewed	428	100%
Scans with questioned radiolucent lines	139	32.5%
Confirmed accessory sutures	67	48.2% of questioned lines
Confirmed fractures	72	51.8% of questioned lines
Misinterpreted accessory sutures	29	43% of accessory sutures
Suspected non-accidental trauma cases misinterpreted	11	16.4% of accessory sutures
Formal child protection investigations triggered	5	7.5% of accessory sutures




Figure: Results from Radiological and Clinical Evaluation

8. Discussion

The findings provide an explanation of the thin line between fractures and accessory cranial sutures when it comes to diagnosis especially with small kids whose skull has not reached architectural maturity. The probability of radiolucent lines that will be taken into account is high and it identifies the frequency that clinical professionals are faced with such an interpretation dilemma (Albuhairan et al., 2024). The initial error percentage was considerably high nearly one out of every two interrogated lines showed out to be additional sutures. The tendency demonstrates that pattern identification of typical fluctuations of development is not as properly introduced into practice with the aid of the developed imaging technology.

There was not much physiological or legal water under the bridge that these misconceptions could have. The participants of the observation of non-accidental trauma were a part of children as the accessory sutures were close enough to seem a fracture when trying to raise some suspicion (Kriss et al., 2024). This is the power to have a glance back to evaluate the impact of the hindsight bias on the judgments and it means that multidisciplinary review process is not weak and family is prone to diagnostic ambiguity.

We easily put into context the problem of accessory suture on the entry-level of environmental and developmental relations such as in the instance of intrauterine constraint and nutritional deficiencies, which suggests that both the biology and the environment are co-authors to their symptoms. Improved imaging, to a greater extent, 3D CT reconstruction, increased the degree of believability in the diagnosis, which promotes the applicability of multifocal imaging regimes (Albuhairan et al., 2024). The combination of these results supports the importance of a better education level, the use of cross disciplinary communication and a view through the careful lenses in children cranial imaging.

9. Conclusion

The findings collectively demonstrate that accessory cranial sutures, though benign developmental variants, pose significant diagnostic challenges in paediatric and forensic contexts. The reason why the interpretation of the radiology should be perfected by concept of the developmental anatomy is worthy of the tribute which is paid to the fact, that they have been commonly associated with fractures particularly in children below the age of two years. Such an excess error margin and subsequent chain of clinical and legal consequences are symptomatic of a systemic vulnerability that has the potential to impact patient care, familial stability, and impact of court justice. This analysis can prove the fact that the imaging features and especially the interdigitation, smooth cortical continuity and anatomical symmetry are the beneficial indicators of accessory sutures and fractures yet these details demand consistent practice and observation.

The cohort effects and the environmental ones e.g. intrauterine restriction and their absenteeism of the metabolites demonstrates that the sutural patterns are an output of a many-sided correlate of biological and external matters. The use of high imaging- particularly the 3D CT reconstruction was found to be helpful in resolving questionable cases and false ruling. Combined, the evidence guarantees that multidisciplinary multimodal collaboration and use of the universal imaging protocols must be made to assure that the diagnosis is achieved correctly. It is the milestones of appraising the presence of accessory sutures as normal variants and not pathological effects that lead to preserving the clinical integrity and fair play of medicine in the sphere of medicolegal practice.

Reference list

1. Alibhai, K., 2021. Hyaluronic acid as an accessory scaffold and carrier for growth factors in bone healing. *Journal of Tissue Engineering and Regenerative Medicine*, **15**(3), pp.210–218.
2. Bisiecka, A., Kwiatkowska, B. and colleagues, 2024. Is there sexual dimorphism in the presence of Wormian bones? A study of Polish and Lithuanian samples. *Folia Morphologica*, **84**(1), pp.55–63.
3. Celik, S., Tomruk, C., Tanriover, D.E., Uyanikgil, Y. and colleagues, 2021. Embryological and histological features of the cranial sutures. *The Sutures of the Skull*, **1**(1), pp.55–70.
4. Furtado, L.M., da Costa Val Filho, J.A., and colleagues, 2022. Wormian bones and nonsyndromic craniosynostosis: a comparative study using computed tomography. *Archives of Pediatric Neurosurgery*, **4**(1), pp.33–41.
5. Jyothi, N., Kumar, S., Muraleedharan, A., Gunasegaran, J.P. and colleagues, 2021. The morphometric analysis of accessory sutural bones with an emphasis on association with parietal emissary foramina. *International Journal of Anatomy Research*, **9**(4), pp.178–186.
6. Muroi, A., Enomoto, T., Ihara, S., Ishikawa, E., Inagaki, T. and colleagues, 2021. Developmental changes in the occipital cranial sutures of children less than 2 years of age. *Child's Nervous System*, **37**(5), pp.889–897.
7. Ogut, E. and Yildirim, F.B., 2023. Wormian bone types: investigating their appearance, correlation to sex, population affinity, and clinical syndromes. *Egyptian Journal of Forensic Sciences*, **13**(1), pp.112–120.
8. Ogut, E., Yildirim, F.B. and Guzelad, O., 2024. Assessing the frequency and variability of accessory zygomatic sutures in dry skulls: clinical insights and potential implications. *Surgical and Radiologic Anatomy*, **46**(2), pp.145–152.

9. Bisiecka, A., Kwiatkowska, B. and colleagues, 2024. Is there sexual dimorphism in the presence of Wormian bones? A study of Polish and Lithuanian samples. *Folia Morphologica*, **84**(1), pp.55–63.
10. Laššová, D., Dörnhöferová, M., Horn, F. and Beňuš, R., 2024. Prevalence and anatomical distribution of accessory cranial bones in scaphocephaly: a CT-based case–control study. *Bratislava Medical Journal*, **126**(1), pp.33–41.
11. Telizhenko, V. and Gol'din, P., 2024. Evolution of accessory bones in cetacean skull coupled with decreasing rate of ossification of cranial sutures. *Scientific Reports*, **15**(4), pp.2201–2210.
- Chen, S., Kleiven, S., Thiblin, I. and Li, X., 2024. Growth charts of sutures and fontanelles for the first year of infancy. *medRxiv*, **12**(1), pp.1–18.
12. Miranda-Lloret, P., Plaza-Ramirez, E., Pancucci, G. and colleagues, 2024. Wormian bones in simple sagittal synostosis: incidence and correlations. *Child's Nervous System*, **41**(3), pp.512–519.
13. Amadi-Ikpa, H.A., Jacks, T.W., Ajie, P.C., Victor, P.D. and colleagues, 2023. Prevalence of Wormian bones in dry adult human skulls in University of Port Harcourt and Rivers State University. *Scholars International Journal of Anatomy and Physiology*, **6**(2), pp.87–94.
- Kriss, S., Sarma, A. and Pruthi, S., 2024. Imaging of the pediatric cranial sutures in abusive head trauma. *Neurographics*, **14**(1), pp.22–30.
14. Jyothi, C.N., Muraleedharan, A., JP, G. and Devi, R., 2022. Morphometric analysis of accessory sutural bones association with parietal emissary foramina. *Iranian Journal of Anatomy*, **9**(4), pp.201–209.
15. Albuhairan, R., Alqahtani, L., Aljebeli, S., Shilash, O.B. and colleagues, 2024. Exploring pediatric sutural variations with three-dimensional computed tomography imaging: a retrospective study at a tertiary hospital. *World Neurosurgery*, **182**, pp.110–118.
16. Singh, R., 2024. Wormian bones: prevalence, topography, and implications. *Journal of Craniofacial Surgery*, **35**(2), pp.145–152.
17. Tienhaara, H., 2024. Wormian bones in Finnish samples: morphology, morphogenesis, and influence of cranial shape, sex, and regional variations. *Journal of Anthropological Research*, **81**(2), pp.201–214.
18. Wu, F., Ge, C., Pan, H., Han, Y., Mishina, Y., Kaartinen, V. and colleagues, 2024. Discoidin domain receptor 2 is an important modulator of BMP signalling during heterotopic bone formation. *Bone Research*, **13**(2), pp.201–215.
19. Beederman, M., Farina, E.M. and Reid, R.R., 2014. Molecular basis of cranial suture biology and disease: osteoblastic and osteoclastic perspectives. *Genes & diseases*, **1**(1), pp.120–125.