



## ‘TO STUDY THE SURGICAL ABDOMINAL SEPSIS (SABS) OF ICU PATIENTS WITH BOWEL PERFORATION PERITONITIS PATIENTS’

Dr. Ravi Pratap Singh<sup>1\*</sup>, Dr. Ashok Kumar Chopda<sup>2</sup>, Dr. Atul Kumar Khare<sup>3</sup>  
Dr. Tanuj Ahirwar<sup>4</sup>

<sup>1\*</sup>Senior Resident, Department Of General Surgery, Dr. Baba Saheb Ambedker Medical College And Hospital Delhi

<sup>2</sup>Senior Resident, Department Of Pediatrics Surgery, Sms Medical College Jaipur

<sup>3</sup>Senior Resident, Department Of Pediatrics Surgery, Sms Medical College Jaipur Orcid Id= 0000-0002-8869-5614

<sup>4</sup>Senior Resident, Department Of General Surgery, Atal Bihari Vajpayee Gmc Vidisha Mp

**\*Corresponding Author:** Dr. Tanuj Ahirwar

<sup>4</sup>Senior Resident, Department Of General Surgery, Atal Bihari Vajpayee Gmc Vidisha Mp

---

### Abstract

**Introduction-** Sepsis is defined as life-threatening organ dysfunction caused by a dysregulated host response to infection. It's the primary cause of death from infection, especially if not recognized and treated promptly. It's a syndrome caused by pathogen factors and host factors with characteristics that evolve over time. What differentiates sepsis from infection is an aberrant or dysregulated host response and the presence of organ dysfunction. The clinical and biological phenotype of sepsis can however be modified by preexisting acute illness, long-standing co-morbidities, medication, and interventions.

**Methods and materials-** This is a prospective cohort study done in tertiary care center in central India “ ” was conducted in Department of Surgery in people's college of medical science and research Centre, Bhopal for two-year study in 50 patients, Surgical Abdominal Sepsis admitted in Surgical intensive care unit (SICU) during study period, for two years periods. Inclusion criteria –All patients belonging to  $\geq 16$  years of age who were admitted in SICU, as a result of perforation peritonitis, requiring exploratory laparotomy, Patients willing to participate in the study after written informed consent. Exclusion criteria- Age  $< 16$  years, Sepsis secondary to trauma, Laparotomies for non-septic indications. After obtaining Ethical clearance from Institute's Ethical Committee, all the patients fulfilling the inclusion criteria were enrolled in the study. Written consent was obtained from all the study participants. The participants were divided in two groups i.e. survivors and non survivors based on their outcome.

**Results-** -Mean age of patients in present study was  $36 \pm 16.14$  years. In present study, 42 (84%) out of 50 patients were males while 8 (16%) were females. All 5 patients who died were males. Most common etiology of SABS was small intestine perforation (36%) followed by gastric perforation (26%). Mean Apache score was maximum in intra-abdominal abscess (105) followed by ischemia (57) and gastric perforation and was minimum in large intestine perforation (29.3). Mean of MAP of

45 patients in survivor group was 82.55 mm Hg while the same in non-survivors was 75 mm Hg. Mechanical ventilation was used in total of 3 patients, out of which 1 survived and 2 patients succumbed to death. The association of mechanical ventilation usage with mortality (chi square test) amongst survivors and non survivors was found to be statistically significant ( $p=0.0008$ ). Deranged Sodium was seen in 11 out of 45 survivors (24.4%) and 2 out of 5(40%) nonsurvivors. The survivors had a mean urine output of 1020 ml and non-survivors had a mean urine output of 350 ml. In case of serum creatinine, mean of survivor group was 0.98 while in non-survivors 1.74. Mean albumin in survivor group was 3.09 mg/dl and in non-survivors was 1.94. In present study, 12 patients had APS score less than 30 with no mortality. 1 out of 31 patients in 30-60 score range died while 4 out of 7 died in case of score  $> 60$ . The mean observed length of stay of all patients was 6.65 days against an expected LOS (length of stay) of 4.6 days. Overall O:E Ratio was 1.44 against O:E of 1.21 in non-survivors with observed and expected LOS of 10.2 and 8.42 days. The difference was statistically significant with  $p$  value = 0.02.

**Conclusion-** From this study we concluded that, results in terms of Length of stay are not accurate and a further workup on the same is required. Although, the results may be improved after removing the limitation of the study which is small sample size.

Hence, a larger study with greater sample size and if possible, involvement of multiple centers is recommended which could help in reaching some conclusive and milestone results.

From this study we evaluate mortality of ICU patients with perforation peritonitis patients, requiring emergency laparotomy, which is essential for all healthcare workers.

**Keywords-** SURGICAL ABDOMINAL SEPSIS (SABS), SICU, emergency laparotomy, perforation peritonitis.

### **Introduction-**

SURGICAL ABDOMINAL SEPSIS (SABS)<sup>[6]</sup> Defined as patients with evidence of preoperative severe sepsis or septic shock with a suspected or known abdominal source of infection requiring laparotomy for source control.

Severe Sepsis is defined as meeting at least one of the following criteria of SIRS with evidence of organ dysfunction.<sup>[7]</sup>

- TEMPERATURE  $> 38^{\circ}\text{C}$  or  $<36^{\circ}\text{C}$
- HR  $> 90/\text{minute}$
- RR  $> 20 / \text{minute}$  or  $\text{PaCO}_2 < 32 \text{ mm Hg}$
- WBC Count  $> 12,000 \text{ cells}/\text{mm}^3$  or  $< 4,000 \text{ cells}/\text{mm}^3$  or  $> 10\%$  immature bands.
- Evidence of Organ dysfunction
- Hypoperfusion [Lactate  $> 1 \text{ mmol/L}$ ]
- Hypotension [SBP  $< 90 \text{ mmHg}$ ]

APACHE (Acute Physiology and Chronic Health Evaluation) score is one of the several intensive care unit (ICU) scoring systems. The death rate of patients admitted to ICU's is much higher than that of other areas.

APACHE has two components: Acute Physiology Score (APS) and the chronic health evaluation, which incorporates the influence of co-morbid conditions (such as diabetes and cirrhosis). Out of these, physiological scoring is aimed at measuring the immediate status of the patient and focuses on the vital organs and their functional evaluation. It is a reflection of long-term disease process. Results of the evaluation can be used to estimate the mortality rate for patients in the ICU and during the hospitalization. <sup>[1]</sup>

### **Aim and objectives-**

This is a prospective cohort study done in tertiary care center in central India ““**To study the SURGICAL ABDOMINAL SEPSIS (SABS) of ICU patients with bowel perforation peritonitis patients**”” was conducted in Department of Surgery in people’s college of medical science and research Centre, Bhopal for two-year study in 50 patients, Surgical Abdominal Sepsis admitted in Surgical intensive care unit (SICU) during study period, for two years periods

### **Inclusion criteria –**

All patients belonging to  $\geq 16$  years of age who were admitted in SICU, patients with perforation peritonitis, requiring exploratory laparotomy. Patients willing to participate in the study after written informed consent.

### **Exclusion criteria-**

- Age < 16 years
- Sepsis secondary to trauma
- Laparotomies for non-septic indications

### **Methodology-**

After obtaining Ethical clearance from Institute’s Ethical Committee, all the patients fulfilling the inclusion criteria were enrolled in the study. Written consent was obtained from all the study participants. Physiological data and Biochemical data were collected at the time of admission or within 24 hours of admission.

The variables used included age, sex, dates of admission, discharge or death, systolic and diastolic blood pressure, body temperature, heart rate, respiratory rate, glucose, blood urea nitrogen, serum sodium, creatinine, blood hematocrit, white blood cells, serum albumin and bilirubin, urine output during the first 24 h of ICU admission, pH, fraction of inspired oxygen, partial pressure of carbon dioxide and partial pressure of oxygen.

Death or discharge and length of stay in ICU were followed up by referring to patients’ medical records. Additionally, APACHE-IV score, Glasgow coma score (GCS), and acute physiology score (APS) were calculated by APACHE IV calculator. (Online tool)

The present study enrolled 50 patients and the participants were divided in two groups i.e. survivors and non survivors based on their outcome.

Data was compiled using MS excel and comparison of APACHE IV score was made between the two groups (survivors and non-survivors) and its predictive efficacy was analysed.

### **Statistical analysis-**

APACHE scores and outcome variables were compared between survivors and non-survivors using Student’s t-test/Fischer exact test wherever applicable.

#### **• Data Analysis was done using SPSS software.**

The present study enrolled 50 patients fulfilling inclusion criteria during the study period. The patients were observed during hospital stay until final outcome i.e. discharge or death. On the basis of their final outcome i.e. survival; patients were divided in two groups: - survivors and non-survivors.

In 2016, the Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3) [8] were released according to which, SIRS (Systemic Inflammatory Response Syndrome) is defined as the presence of Two or more of:

- Temperature  $>38^{\circ}\text{C}$  or  $< 36^{\circ}\text{C}$

- Heart rate  $>90/\text{min}$
- Respiratory rate  $>20/\text{min}$  or  $\text{PaCO}_2 < 32 \text{ mm hg}$ .
- WBC count  $> 12\,000/\text{mm}^3$  or  $< 4000/\text{mm}^3$  or  $> 10\%$  immature bands

Sepsis is defined as life-threatening organ dysfunction caused by a dysregulated host response to infection. It's the primary cause of death from infection, especially if not recognized and treated promptly. It's a syndrome caused by pathogen factors and host factors (e.g, sex, race and other genetic determinants, age, co-morbidities, environment) with characteristics that evolve over time. What differentiates sepsis from infection is an aberrant or dysregulated host response and the presence of organ dysfunction. The clinical and biological phenotype of sepsis can however be modified by preexisting acute illness, long-standing co- morbidities, medication, and interventions.

Sepsis-induced organ dysfunction may be occult; therefore, its presence should be considered in any patient presenting with infection. Patients with suspected infection who are likely to have a prolonged ICU stay or to die in the hospital can be promptly identified at the bedside with qSOFA, ie, alteration in mental status, systolic blood pressure  $<100 \text{ mm Hg}$ , or respiratory rate  $\geq 22/\text{min}$ .

Septic shock is a further subset of sepsis in which underlying circulatory and cellular/metabolic abnormalities are profound enough to substantially increase mortality. Patients with septic shock can be identified with a clinical construct of sepsis with persisting hypotension requiring vasopressors to maintain  $\text{MAP} \geq 65 \text{ mm Hg}$  and having a serum lactate level  $>2 \text{ mmol/L}$  ( $18 \text{ mg/dL}$ ) despite adequate volume resuscitation.

### **Fig:1 Clinical features for Identifying Patients with Sepsis and Septic Shock**

Following pulmonary sepsis, abdominal sepsis is the second most common form of sepsis requiring intensive care unit (ICU) management. In a nation-wide survey of ICU-treated adults with severe sepsis in Finland, an intra-abdominal source was found in 32% of patients and was associated with a hospital mortality rate of 30%.<sup>[10]</sup>

### **Prediction Scores-**

Scoring systems have been developed in response to an increasing emphasis on the evaluation and monitoring of health services. These systems enable comparative audit and evaluative research of intensive care. The ideal components of a scoring system are data collected during the course of routine patient management that are easily measured, objective, and reproducible. Scoring systems, developed in the 1980s are applicable to heterogeneous groups of critically ill patients.

The evaluation of severity of illness in the critically ill patient is made through the use of severity scores and prognostic models. Severity scores are instruments that aim at stratifying patients based on the severity of illness, assigning to each patient an increasing score as their severity of illness increases. Prognostic models, apart from their ability to stratify patients according to their severity, predict a certain outcome (usually the vital status at hospital discharge) based on a given set of prognostic variables and a certain modeling equation.<sup>[20]</sup>

Previously known as Sepsis related Organ failure assessment, The Sequential Organ Failure Assessment (SOFA) score was developed to quantify the severity of patients' illness, based on the degree of organ dysfunction, was introduced in 1994 and further revised in 1996.

The SOFA score is composed of scores from six organ systems, each graded from 0 to 4 points according to the degree of dysfunction, giving a possible score of 0 to 24. The SOFA scoring system takes into account the time course of a patient's condition during the entire ICU stay.

Organ dysfunction can be identified as an acute change in total SOFA score  $\geq 2$  points consequent to the infection. The baseline SOFA score can be assumed to be zero in patients not known to have

preexisting organ dysfunction.

ASOFA score  $\geq 2$  reflects an overall mortality risk of approximately 10% in a general hospital population with suspected infection. [27]

**Observation and results-**

-In present study, 42 (84%) out of 50 patients were males while 8 (16%) were females. All 5 patients who died were males.

-Mean age of patients in present study was  $36 \pm 16.14$  years. Out of total 50 patients, 7 patients belonged to less than 20 years of age, with no mortality. 33 patients belonged to 20-50 years of age group with 3 deaths and 10 patients belonged to more than 50 years with 2 deaths.

-Most common etiology of SABS was small intestine perforation (36%) followed by gastric perforation (26%), Small intestinal obstruction (14%), appendicular perforations (8%), large bowel perforations (6%), large bowel obstructions (4%) and intra-abdominal abscess (1%). Mean Apache score was maximum in intra-abdominal abscess (105) followed by ischemia (57) and gastric perforation and was minimum in large intestine perforation (29.3).

**Table 1 Etiological Breakup of Diseases with Mortality**

Etiology	No of Patients	Survivors	Non-survivors	Mean ApacheScore
Large IntestinePerforation	3(6%)	3	0	29.3
Small IntestinePerforation	18(36%)	18 (100%)	0	43
Gastric Perforation	13(26%)	10(76.92%)	3(23.07%)	50.4
Abscess (Intra-abdominal)	1(2%)	0	1(100%)	105
AppendicularPerforation	4(8%)	4(100%)	0	43.5
Large IntestineObstruction	2(4%)	2(100%)	0	34
Small IntestineObstruction	7(14%)	7(100%)	0	35
Ischemia/Infarction	2(4%)	1(50%)	1(50%)	57

-Mean of MAP of 45 patients in survivor group was 82.55 mm Hg while the same in non-survivors was 75 mm Hg. Test of significance could not be applied as the data in the two groups was less.

**Table 2 MAP values in survivors and non survivors**

	No of Patients	MAP (mm Hg)
Survivors	45	82.55
Non-Survivors	5	75

**Table 3 Use of Mechanical Ventilation in Survivors and Non- Survivors**

Mechanical VentilationUsed	Survivors	Non-Survivors
Yes	1	2
No	44	3
P = 0.0008		

-Mechanical ventilation was used in total of 3 patients, out of which 1 survived and 2 patients succumbed to death. The association of mechanical ventilation usage with mortality (chi square test) amongst survivors and non survivors was found to be statistically significant (p=0.0008).

-Deranged Sodium was seen in 11 out of 45 survivors (24.4%) and 2 out of 5(40%)nonsurvivors. The association of mortality with deranged sodium level using chi square test was found to be statistically insignificant in present study (p=0.45)

### Apache Score-

There was no mortality in 43 patients with APACHE Score <60. 60-80 score group had 2 patients with no mortality and 5 patients belonged to score > 80 with 100% mortality. For the ease of analysis, Apache score was divided into ≤80 and >80. Test of significance (Fischer exact test) showed statistically significant association between mortality and Apache score (p<0.0001). Mean APACHE score of Survivor group was 39.0 while that of non-survivor group was 86.20 and overall mean of all patients was 44.46.

**Table 8 (a)-Division of cases on basis of Apache score**

Apache Score	Total Patients	Mortality
<60	43	0
60-80	2	0
>80	5	5
<b>P &lt; 0.0001</b>		

**Table 8 (b)- Mean APACHE Scores of both groups**

Group	Mean Apache Score
Survivors (n=45)	39.0
Non-Survivors (n=5)	86.20
Overall (n=50)	44.46

### APS Score-

-In present study, 12 patients had APS score less than 30 with no mortality. 1 out of 31 patients in 30-60 score range died while 4 out of 7 died in case of score > 60.

**Table 9 Division of cases on basis of APS score**

APS Score	Total Patients	Mortality
<30	12	0
30-60	31	1
>60	7	4

### Estimated Mortality Rates

-A receiver operating characteristic (ROC) curve, comparative analysis of the mortality predictions with APACHE IV is done and shows AUC =0.911 and p <0.001. Thus, showing the high predictive efficacy of APACHE IV score.

### Length of Stay( LOS )-

-The mean observed length of stay of all patients was 6.65 days against an expected LOS (length of stay) of 4.6 days. Overall O:E Ratio was 1.44 against O:E of 1.21 in non-survivors with observed and expected LOS of 10.2 and 8.42 days. The difference was statistically significant with p value = 0.02

**Table 10 Comparison of Observed and Expected LOS**

No of Patients	Observed LOS(Days)	Expected LOS(Days)	O: E Ratio
50	6.65	4.6	1.44
5 (non-Survivors)	10.2	8.42	1.21
<b>P value 0.0270</b>			

### Discussion-

The present study entitled “**Evaluation of Predictive Efficacy of Apache IV Score in Surgical Abdominal Sepsis**” was conducted on 50 patients admitted in S.I.C.U. diagnosed with Surgical Abdominal Sepsis requiring exploratory laparotomy. Mean age of patients in our study was 36±16.14 years. 33 patients belonged to 20-50 years of age followed by 10 patients of >50 years age and 7 patients of <20 years of age. Whereas mortality was observed in 2 and 3 patients belonging to >50

years and 20-50 years of age group respectively. There was an age- wise increase in the rate of mortality as in the results above but the relation was not found to be statistically significant. Similar findings were seen in the study by **Moses S et al.** who also had similar results.<sup>[1]</sup> In another study by **Chan T et al**, the mean age of study participants was 62.8±15.2 years. They found a significant difference in age between survivors (60.7 years, 95% CI 58.3, 63.2) and non-survivors (67.9 years, 95% CI 64.1, 71.6).<sup>[32]</sup>

Majority of participants in present study were males that is 84% while only 16% participants were females. Mortality was observed in 5 males in our study. In a study by **Ghorbani M et al** in Iran, study population comprised of 53.9% males and 46.1% females.<sup>[33]</sup> In another study by **Shoukat H et al** 55.5% and 44.5% participants were males and females respectively. Out of 86 male patients, mortality occurred in 41 (47.67%) patients and did not occur in 45 (52.33%) patients, while 69 patients were females in which mortality occurred in 31 (44.92%) and not in 38 (55.08%) patients.<sup>[38]</sup> Most common etiology observed in present study was small intestine perforation (36%), followed by gastric perforation (26%) and Small intestinal obstruction (14%).

Similar disease patterns were found in the study by **Vishwani et al**<sup>[39]</sup> and **Shoukat H et al**<sup>[38]</sup>, both of which included patients of sepsis undergoing exploratory laparotomy.

In present study, based on mortality, patients were divided into 2 groups, survivor and non survivor. Mean of MAP of 45 patients in survivor group was 82.55 mm Hg while the same in non-survivors was 75 mm Hg. Not much studies have considered mean arterial pressure. The abdominal perfusion pressure (mean arterial pressure– intra- abdominal pressure) has been shown to correlate with survival when maintained at levels greater than 50 mm Hg to 60 mm Hg, but level I evidence examining this as an end point of resuscitation has yet to be obtained.<sup>[40]</sup>

In present study, mechanical ventilation was used in 3 patients. The association of mechanical ventilation usage with mortality amongst survivors and non survivors was found to be statistically significant ( $p=0.0008$ ). **Martin ND et al** in their study could not assess the association of mechanical ventilation and mortality, since mechanical ventilation was required in all patients.<sup>[41]</sup>

The present study observed insignificant association of mortality with deranged sodium level. The mean urine output among survivors was 1020 ml and that in non- survivors was 350 ml. In case of serum creatinine, mean of survivor group was 0.98 while in non-survivors is 1.74. Mean albumin in survivor group was 3.09 mg/dl and in non-survivors was 1.94. Test of significance could not be applied as the data was less. The development of Acute Kidney injury is common following laparotomy, reaches greatest prevalence 48h after initial laparotomy, and is an independent predictor of increased mortality.<sup>[42]</sup>

In present study it was observed, higher the Apache score, worst is the prognosis and more the mortality. Test of significance showed statistically significant association between mortality and Apache score ( $p<0.0001$ ). **Ghorbani M et al** also observed significant difference in mean Apache score amongst survivors and non survivors ( $p<0.001$ ).<sup>[33]</sup> **El-Naggar TA et al** in their study observed APACHE IV scores were significantly higher between dead than alive patients on admission and after 48 h, but were not able to predict death in ICU.<sup>[43]</sup> **Chan T et al** suggested APACHE-IV crudely distinguished between survivors and non-survivors.<sup>[32]</sup>

Also present study observed higher the Apache score, worst the prognosis and more the mortality. These findings were similar to study by **El-Naggar TA et al**<sup>[43]</sup> and **Saleh A et al.**<sup>[44]</sup> **Yamin S et al** in their study observed mean APACHE IV score of survivors to be 54.55, while mean APACHE IV score of non survivor was 85.07 which was significantly higher. 63.9% patients had APACHE IV score <60, out of these 14.8% didn't survive. 27.8% patients had APACHE IV score >81 out of these



65.8% didn't survive ( $p < 0.001$ ). The 62.1% of overall population show the same outcome as predicted by APACHE IV ( $p = 0.61$ ).<sup>[45]</sup>

The ROC curve showed the results to be pretty accurate with an AUC = 0.91 and  $p < 0.01$ . AUC was 0.93 in a study in Turkey by **Ayazoglu TA et al.**<sup>[46]</sup> In a study in South Korea also, AUC was 0.80 and significantly better than its predecessors.<sup>[47]</sup> **Zimmerman JE et al** concluded APACHE IV had good discrimination (area under the receiver operating characteristic curve = 0.88) and calibration (Hosmer-Lemeshow C statistic = 16.9,  $p = .08$ ). For 90% of 116 ICU admission diagnoses, the ratio of observed to predicted mortality was not significantly different from 1.0.<sup>[28]</sup> **Kuzniewicz MW et al** also found similar results. They observed APACHE IV had the best discrimination (area under the receiver operating characteristic curve [AUC], 0.892) compared to MPM(0) III (AUC, 0.809), and SAPS II (AUC, 0.873;  $p < 0.001$ ).

The models differed substantially in data abstraction times, as follows: MPM(0)III, 11.1 min (95% confidence interval [CI], 8.7 to 13.4); SAPS II, 19.6 min (95% CI, 17.0 to 22.2); and APACHE IV, 37.3 min (95% CI, 28.0 to 46.6).<sup>[48]</sup>

In terms of Length of stay (LOS), Mean of all patients was 6.65 days against an expected LOS of 4.6 days. Overall O:E (Observed LOS : Expected LOS) Ratio was 1.44 against, O:E of 1.21 in non-survivors with observed and expected LOS of 10.2 and 8.42 days. The results were pretty accurate with higher accuracy in non-survivor than the survivor group. The slightly higher prediction of Length of stay in survivor group by APACHE IV may be because of a conservative approach to shift and discharge patients in the institute because of most patients were from the rural and low socio-economic group and hence the patient factors have to be kept in mind.<sup>[49]</sup>

In a pilot study predicting ICU length of stay using APACHE-IV in severe sepsis patients by **Chattopadhyay and Chatterjee in Cleveland**, Ohio, similar results was found with a significant over-prediction of length of stay.<sup>[50]</sup>

Lack of inpatient bed availability is a major contributor to emergency department (ED) crowding. Important reductions in LOS at ED (and others) and smoothening of ambulance diversion occur in urban academic medical center after an increase in adult ICU beds. A recent study demonstrated that the most notable change after ICU expansion was a decrease in time spent on ambulance diversion. Increasing ICU beds appears to have shortened ED LOS for ICU patients but has less effect on other admitted patients and apparently, no effect on patients discharged home. In emergency care a perfect balance between given resources and demand is much more difficult to achieve and maintain. In reality, certain key resources, e.g. ICU beds, are often over-utilized.<sup>[18]</sup>

### **Conclusion-**

From this study we concluded that, results in terms of Length of stay are not accurate and a further workup on the same is required. Although, the results may be improved after removing the limitation of the study which is small sample size.

Hence, a larger study with greater sample size and if possible, involvement of multiple centers is recommended which could help in reaching some conclusive and milestone results.

### **References**

1. Moses S, Gautam A, Shukla S, Mathur RK. Evaluation of predictive efficacy of APACHE IV score in abdominal trauma patients. JOURNAL OF EVOLUTION OF MEDICAL AND DENTAL SCIENCES-JEMDS. 2015 ;4(28):4834-43.
2. Knaus WA, Draper EA, Wagner DP, Zimmerman JE. APACHE II: a severity of disease classification system. Critical care medicine. 1985;13(10):818-29.



3. Knaus WA, Wagner DP, Draper EA, Zimmerman JE, Bergner M, Bastos PG, Sirio CA, Murphy DJ, Lotring T, Damiano A, Harrell Jr FE. The APACHE III prognostic system: risk prediction of hospital mortality for critically III hospitalized adults. *Chest*. 1991;100(6):1619-36.
4. Knaus WA, Zimmerman JE, Wagner DP, Draper EA, Lawrence DE. APACHE-acute physiology and chronic health evaluation: a physiologically based classification system. *Critical care medicine*. 1981;9(8):591-7.
5. Zimmerman JE, Kramer AA, McNair DS, Malila FM, Shaffer VL. Intensive care unit length of stay: Benchmarking based on Acute Physiology and Chronic Health Evaluation (APACHE) IV. *Critical care medicine*. 2006;34(10):2517-29.
6. Hanisch E, Brause R, Paetz J, Arlt B. Review of a large clinical series: predicting death for patients with abdominal septic shock. *Journal of intensive care medicine*. 2011;26(1):27-33.
7. Singer M, Deutschman CS, Seymour CW, Shankar-Hari M, Annane D, Bauer M, Bellomo R, Bernard GR, Chiche JD, Coopersmith CM, Hotchkiss RS. The third international consensus definitions for sepsis and septic shock (Sepsis-3). *Jama*. 2016;315(8):801-10.
8. Lipinska-Gediga M. Coagulopathy in sepsis—a new look at an old problem. *Anaesthesiology intensive therapy*. 2016;48(5):352-9.
9. Marik PE, Taeb AM. SIRS, qSOFA and new sepsis definition. *Journal of thoracic disease*. 2017;9(4):943.
10. Friedman G, Silva E, Vincent JL. Has the mortality of septic shock changed with time?. *Critical care medicine*. 1998;26(12):2078-86.
11. Leppäniemi A, Kimball EJ, Malbrain ML, Balogh ZJ, De Waele JJ. Management of abdominal sepsis—a paradigm shift?. *Anaesthesiology intensive therapy*. 2015;47(4):400-8.
12. Pieracci FM, Barie PS. Management of severe sepsis of abdominal origin. *Scandinavian journal of surgery*. 2007;96(3):184-96.
13. Iwashyna TJ, Cooke CR, Wunsch H, Kahn JM. Population burden of long-term survivorship after severe sepsis in older Americans. *Journal of the American Geriatrics Society*. 2012;60(6):1070-7.
14. Dahhan T, Jamil M, Al-Tarifi A, Abouchala N, Kherallah M. Validation of the APACHE IV scoring system in patients with severe sepsis and comparison with the APACHE II system. *Critical Care*. 2009;13(1):P511.
15. Le Gall JR, Lemeshow S, Saulnier F. A new simplified acute physiology score (SAPS II) based on a European/North American multicenter study. *Jama*. 1993;270(24):2957-63.
16. Kirkpatrick AW, Coccolini F, Ansaloni L, Roberts DJ, Tolonen M, McKee JL, Leppaniemi A, Faris P, Doig CJ, Catena F, Fabian T. Closed Or Open after Source Control Laparotomy for Severe Complicated Intra-Abdominal Sepsis (the COOL trial): study protocol for a randomized controlled trial. *World Journal of Emergency Surgery*. 2018;13(1):26.
17. Papavramidis TS, Marinis AD, Pliakos I, Kesisoglou I, Papavramidou N. Abdominal compartment syndrome—Intra-abdominal hypertension: Defining, diagnosing, and managing. *Journal of emergencies, trauma and shock*. 2011;4(2):279.
18. Strik C, Stommel MW, Schipper LJ, van Goor H, ten Broek RP. Risk factors for future repeat abdominal surgery. *Langenbeck's archives of surgery*. 2016;401(6):829-37.
19. Mehrez A, Gafni A. Preference Based Outcome Measures for Economic Evaluation of Drug Interventions. *Pharmacoeconomics*. 1992;1(5):338-45.
20. Gunning K, Rowan K. Outcome data and scoring systems. *Bmj*. 1999 ;319(7204):241-4.
21. Rapsang AG, Shyam DC. Scoring systems in the intensive care unit: a compendium. *Indian journal of critical care medicine: peer-reviewed, official publication of Indian Society of Critical Care Medicine*. 2014;18(4):220.
22. Vincent JL, Moreno R. Clinical review: scoring systems in the critically ill. *Critical care*. 2010;14(2):207.
23. Becker RB, Zimmerman JE. ICU scoring systems allow prediction of patient outcomes and comparison of ICU performance. *Critical care clinics*. 1996 ;12(3):503-14.

24. Vincent JL, Marshall JC, Namendys-Silva SA, François B, Martin-Loeches I, Lipman J, Reinhart K, Antonelli M, Pickkers P, Njimi H, Jimenez E. Assessment of the worldwide burden of critical illness: the intensive care over nations (ICON) audit. *lancet Respiratory medicine*. 2014;2(5):380-6.
25. Bota DP, Melot C, Ferreira FL, Ba VN, Vincent JL. The multiple organ dysfunction score (MODS) versus the sequential organ failure assessment (SOFA) score in outcome prediction. *Intensive care medicine*. 2002;28(11):1619-24.
26. Vincent JL, Moreno R, Takala J, Willatts S, De Mendonça A, Bruining H, Reinhart CK, Suter P, Thijs LG. The SOFA (Sepsis-related Organ Failure Assessment) score to describe organ dysfunction/failure.
27. Simalti AK, Ghuliani R, Nair BT. Update on Changing Terminologies in Sepsis. *Journal of Nepal Paediatric Society*. 2016;36(2):178-83.
28. Zimmerman JE, Kramer AA, McNair DS, Malila FM. Acute Physiology and Chronic Health Evaluation (APACHE) IV: hospital mortality assessment for today's critically ill patients. *Critical care medicine*. 2006;34(5):1297-310.
29. Bhattacharyya M, Todi S. APACHE IV: benchmarking in an Indian ICU. *Critical Care*. 2009;13(1):P510
30. Anderson ID, Fearon KC, Grant IS. Laparotomy for abdominal sepsis in the critically ill. *British journal of surgery*. 1996;83(4):535-9.
31. McLauchlan GJ, Anderson ID, Grant IS, Fearon KC. Outcome of patients with abdominal sepsis treated in an intensive care unit. *British journal of surgery*. 1995;82(4):524-9.
32. Chan T, Bleszynski MS, Buczkowski AK. Evaluation of APACHE-IV predictive scoring in surgical abdominal sepsis: a retrospective cohort study. *JCDR*. 2016;10(3):16.
33. Ghorbani M, Ghaem H, Rezaianzadeh A, Shayan Z, Zand F, Nikandish R. A study on the efficacy of APACHE-IV for predicting mortality and length of stay in an intensive care unit in Iran. *IJS*. 2017;6(2).
34. Orsini J, Blaak C, Yeh A, Fonseca X, Helm T, Butala A, Morante J. Triage of patients consulted for ICU admission during times of ICU-bed shortage. *Journal of clinical medicine research*. 2014;6(6):463.
35. Knaus WA, Draper EA, Wagner DP, Zimmerman JE. An evaluation of outcome from intensive care in major medical centers. *Annals of Internal Medicine*. 1986;104(3):410-8.
36. VijayGanapathy S, Karthikeyan VS, Sreenivas J, Mallya A, Keshavamurthy R. Validation of APACHE II scoring system at 24 hours after admission as a prognostic tool in urosepsis: A prospective observational study. *Investigative and clinical urology*. 2017;58(6):453-9.
37. Sadaka F, EthmaneAbouElMaali C, Cytron MA, Fowler K, Javaux VM, O'Brien J. Predicting Mortality of Patients With Sepsis: A Comparison of APACHE II and APACHE III Scoring Systems. *Journal of clinical medicine research*. 2017;9(11):907.
38. Shoukat H, Muhammad Y, Gondal KM, Aslam I. Mortality prediction in patients admitted in surgical intensive care unit by using APACHE IV. *J Coll Physicians Surg Pak*. 2016 ;26(11):877-0.
39. Vishwani A, Gaikwad VV, Kulkarni RM, Murchite S. Efficacy of POSSUM Scoring System in Predicting Mortality and Morbidity in Patients of Peritonitis Undergoing Laparotomy. *Int J Sci Stud*. 2014;2(4):29-36.
40. Chabot E, Nirula R. Open abdomen critical care management principles: resuscitation, fluid balance, nutrition, and ventilator management. *Trauma surgery & acute care open*. 2017;2(1):e000063.
41. Martin ND, Patel SP, Chreiman K, Pascual JL, Braslow B, Reilly PM, Kaplan LJ. Emergency Laparotomy in the Critically Ill: Futility at the Bedside. *Critical care research and practice*. 2018.
42. Loftus TJ, Bihorac A, Ozrazgat-Baslanti T, Jordan JR, Croft CA, Smith RS, Efron PA, Moore FA, Mohr AM, Brakenridge SC. Acute kidney injury following exploratory laparotomy and temporary abdominal closure. *Shock*. 2017;48(1):5-10.
43. El-Naggar TA, Raafat RH, Mohamed SA. Validity of three scoring systems in assessing the

- severity and outcome in Al-Abbassia Chest Hospital Respiratory Intensive Care Unit patients. *Egyptian Journal of Bronchology*. 2018;12(2):208.
44. Saleh A, Ahmed M, Sultan I, Abdel-Lateif A. Comparison of the mortality prediction of different ICU scoring systems (APACHE II and III, SAPS II, and SOFA) in a single-center ICU subpopulation with acute respiratory distress syndrome. *Egyptian journal of chest diseases and tuberculosis*. 2015;64(4):843-8.
  45. Yamin S, VASWANI AK, AFREEDI M. Predictive efficacy of APACHE IV at ICU. s of CHK. *Pakistan Journal of Chest Medicine*. 2011;17(1).
  46. Ayazoglu TA. A comparison of APACHE II and APACHE IV scoring systems in predicting outcome in patients admitted with stroke to an intensive care unit. *Anaesthesia, Pain & Intensive Care*. 2011;15(1).
  47. Lee H, Shon YJ, Kim H, Paik H, Park HP. Validation of the APACHE IV model and its comparison with the APACHE II, SAPS 3, and Korean SAPS 3 models for the prediction of hospital mortality in a Korean surgical intensive care unit. *Korean journal of anesthesiology*. 2014;67(2):115-22.
  48. Kuzniewicz MW, Vasilevskis EE, Lane R, Dean ML, Trivedi NG, Rennie DJ, Clay T, Kotler PL, Dudley RA. Variation in ICU risk-adjusted mortality: impact of methods of assessment and potential confounders. *Chest*. 2008 ;133(6):1319-27.
  49. Chelluri L, Grenvik A, Silverman M. Intensive care for critically ill elderly: mortality, costs, and quality of life: review of the literature. *Archives of Internal Medicine*. 1995;155(10):1013-22.
  50. Chattopadhyay A, Chatterjee S. Predicting ICU length of stay using APACHE-IV in persons with severe sepsis—a pilot study. *Journal of Epidemiological Research*. 2015