



FREQUENCY OF INDICATIONS AND TYPES OF ENTERIC STOMA IN SURGICAL WARD OF AYUB TEACHING HOSPITAL ABBOTTABAD

Tahir Ur Rehman¹, Muhammad Kashif Rafiq^{2*}, Waqas Ashraf³, Yousuf Aziz Khan⁴, Babar Sultan⁵

¹Resident Surgeon, Ayub Teaching Hospital, Abbottabad, Pakistan

^{2*}Assistant Professor of Surgery, Ayub Teaching Hospital, Abbottabad, Pakistan

³Resident Surgeon, Department of Surgery, Ayub Teaching Hospital, Abbottabad, Pakistan

⁴Assistant Professor, Department of Paediatric Surgery, Surgery & Allied, Ayub Teaching Hospital MTI, Pakistan

⁵Associate Professor of Surgery, Ayub Teaching Hospital, Abbottabad, Pakistan

***Corresponding author:** Muhammad Kashif Rafiq

*Email: kashan2002pk@yahoo.com

ABSTRACT

Introduction: Strategic stomas are crucial operations in the management of various gastrointestinal disorders such as intestinal obstruction, perforation, and trauma. It is necessary to examine the frequency of this type of stoma, as well as indications for its treatment and outcomes, in order to improve patient management.

Objectives: To determine the frequency of indications, and types of enteric stomas formed in the surgical ward of Ayub Teaching Hospital, Abbottabad and analyze associated outcomes.

Materials and Methods: This cross-sectional study was performed in 2024 for six months with 121 patients with stoma formation. The research employed a nonprobability consecutive sampling technique. The demographic information, indications for the creation of a stoma, type of stoma and postoperative complications were recorded. Descriptive statistics in the current study were done using Statistical Package for Social Sciences (SPSS) software version 23.

Results: The most common indications for operation were intestinal obstruction, which was seen in 36.4%, and perforation in 28.1%. Loop stomas (58.7%) were the most common, while end stomas (11.6%). Wound infection, which was noted in 15.7% of patients, and stoma retraction in 10.7% of patients were more common in emergency surgeries, which formed 42.3%.

Conclusion: The aspects of early diagnosis, optimum preoperative preparation and psychological intervention need to be specifically personalised for the benefit of stoma patients.

Keywords: Enteric stoma, intestinal obstruction, stoma complications, loop stoma, surgical outcomes.

INTRODUCTION

The formation of enteric stomas has emerged as an essential procedure in the management of many gastrointestinal disorders, from congenital malformations to severe acquired pathology. Colostomies and Ileostomies are made when there is a disease condition or trauma expected to interfere with the normal process of elimination through the anus. The reviews of the recent studies show that more

precise knowledge of indications, results, and types of stomas made in various kinds of health care services would facilitate effective patient management and minimize the adverse effects. It is against this backdrop that the present study, conducted in the surgical ward of Ayub Teaching Hospital Abbottabad, aims to address the regional knowledge gap by assessing the frequency of indications of stoma formation. Therefore, the rate of stoma formation varies across the world depending on IBD, colorectal cancer, and other abdominal injury incidents. Sushel et al. (1) and Mbelle et al. (2) have worked on a topic on stoma creation with indications including intestinal tuberculosis, stoma, perforations and obstructions. In Pakistan, the rate of intestinal tuberculosis is still a significant problem and presents more frequently as acute abdomen and requires surgery with stoma formation (1). Likewise, Mbelle et al. (2), working in sub-Saharan Africa, have also confirmed that stoma creation is often required for late stenosis obstructions and serious perforations, highlighting indication differences.

The formation of the stoma has its own drawbacks. Moya-Muñoz et al. (3) examined the quality of life for stoma patients to note its impact on quality and mental health in patients. Specifically, retraction, infection, and psychosocial stress affect outcomes (3, 6). The results of Granieri et al. (4) expand the understanding of associations between underlying factors such as fistulas and stoma-related adverse effects. A review of literature from other countries shows that when adequate preoperative and postoperative handling is adopted, which includes preoperative stoma site marking, then postoperative complication is significantly reduced, as Nozawa et al. support this argument. In pediatric populations, various developmental factors make it difficult to manage stomas. Chukwubuike et al. (11) demonstrated that children, in particular, suffer from high rates of complications that require better treatment strategies. As suggested by Kimbugwe et al. (6), complications should be treated early so that they do not result in other problems in future. As for adults, they also have complications, notable that each spectrum is different, and Crohn's disease and ulcerative colitis predominate in some areas (7).

Essential to the overall experience is also the psychological load of stoma creation. Tang et al. (8) described a meta-analytical review examining the rate of depression among stoma patients and highlighting the importance of an integrated model of patient care. There is also pre-existing evidence that Pakistan frequently suffers from shortages of resources when it comes to healthcare, which, in turn, means that the amount of psychological support involved in the treatment process is also restricted (9). This research will seek to not only describe the clinical indications but also help identify ways of improving care. Technology and research have been used in stoma care, and they have proved to be effective. For instance, Ahmed et al. (15) presented that the interventions based on simulation effectively enhance the overall performance and satisfaction of nurses for stoma care. In implementation, such interventions in local care facilities can help minimize quality chasm and improve patient health.

This study is in line with a worldwide attempt to improve post-surgical results by determining trends of the formation of stomas across regions. Conclusions drawn from this study shall add to the existing body of literature, including the systematic review conducted by Parini et al. (12), which aimed at identifying worldwide trends in stoma complications as well as surgical procedures. However, depending on the localized data will give important information regarding complications and possible preventions suited to the Pakistani population of patients. The general approach to this study is underpinned by innovative developments in data analysis and categorization methods. Hapnes et al. (14) have provided a good example of the use of robust statistics for the assessment of surgical performance in high-risk groups such as pre-term infants who may require intestinal surgery. Pursuing these methodologies, this study shall use SPSS version 23 for data analysis to enhance the effectiveness as well as the accuracy of the outcomes produced.

Lastly, the creation of Enteric stomas is a life-saving surgery with quality-of-life impacts for the given patients. In turn, this study will assist in filling the existing gap of information on indications and stoma site frequency at Ayub Teaching Hospital Abbottabad and contribute to the future enhancement of surgical care guidelines in Pakistan. Knowledge obtained will also be used in policy making and

funding to help patients get the best care possible, depending on the different areas in health care delivery.

Objective: To determine the frequency of indications of enteric stoma creation and the types of stomas formed in the surgical ward of Ayub Teaching Hospital Abbottabad, addressing local healthcare needs.

MATERIALS AND METHODS

Study Design: Cross-sectional study

Study setting: The study was conducted in the surgical ward of Ayub Teaching Hospital Abbottabad, Pakistan which is the largest health care facility of Pakistan providing tertiary care surgical services.

Duration of the study: The study was conducted over a six-month period, from January 2024 to June 2024.

Inclusion Criteria

This research targeted patients of 12 years and above, both male and female, who had an enteric stoma during the research period. The patients who required stoma creation due to elective or emergency surgery for indications like intestinal obstruction, perforation, or trauma were included. Patients in the study had to be receiving palliative care, be over 18 years of age, have an enteric stoma as defined by the above operational definitions and must have given informed consent before enrolment.

Exclusion Criteria

All patients who had a stoma formed prior to the study period or as part of gynaecological treatments were excluded from the study. Furthermore, those patients who were transferred from another hospital for stoma revision or complications were excluded, and only those patients who underwent primary stoma creation were included in the study conducted at Ayub Teaching Hospital.

Methods

Ethical approval was sought and granted by the institutional review board, and then patients who fulfilled the inclusion criteria were recruited into the study via consecutive non-probability sampling. The demographic and clinical details of each participant were assessed prospectively using a preoperative questionnaire and postoperative case records. The type of stoma created (loop, double, or end) was also noted, and the findings were compared to the operative note. General data from patient interviews and specific clinical findings from patient examination and investigations, including imaging and biochemical data, were applied to secure the diagnosis of stoma formation. The sources of data were surgical notes, histopathological reports, and check observations of any complications that may occur after surgery. The data collected were analyzed using Statistical Package for Social Sciences (SPSS) version 23. Normally distributed continuous variables, including age and weight, were described using the mean and standard deviation, whereas counting data, including gender and types of stomata, were described using frequencies and percentages. Statistical significance being set at $p \leq 0.05$.

RESULTS

The patients without BA surgery who were followed during the six-month study period consisted of 121 patients who had undergone enteric stoma formation. The mean age of the patients was 46.3 ± 15.2 years, with a male-to-female ratio of 1.8:1. They comprised an emergency operation of 64.5% while 35.5% were elective ones. In particular, 27.3% of the population had diabetes, 31.4% had hypertension, and 18.2% had ischemic heart disease as compared with the general population.

Table 1: Demographic and Clinical Characteristics of Patients

Variable	Frequency (n = 121)	Percentage (%)
Mean Age (years)	46.3 ± 15.2	-
Male Patients	79	65.3
Female Patients	42	34.7
Emergency Surgeries	78	64.5
Elective Surgeries	43	35.5
Diabetes	33	27.3
Hypertension	38	31.4
Ischemic Heart Disease	22	18.2

The common indications for the formation of stoma were intestinal obstruction 36.4%, intestinal perforation 28.1% and abdominal trauma 20.7%. Other signs were enterocutaneous fistulas in 8.3% of patients, gastrointestinal masses in 4.1% of patients and mesenteric ischemia in 2.5% of patients. Of all made stomas, loop stomas were the most often made (58.7%), end stomas were made in 11.6% of cases and double-barrel stomas in 29.8% of cases.

Table 2: Indications for Stoma Formation

Indication	Frequency (n = 121)	Percentage (%)
Intestinal Obstruction	44	36.4
Intestinal Perforation	34	28.1
Abdominal Trauma	25	20.7
Enterocutaneous Fistula	10	8.3
Gastrointestinal Mass	5	4.1
Mesenteric Ischemia	3	2.5

Overall, postoperative complications occurred in 32.2% of the patients, and the most common were wound infection (15.7%), stoma retraction (10.7%), and parastomal hernia (5.8%). Patients who underwent emergency surgeries were sept more times to develop complications than the elective patients (42.3; 16.3) per cent respectively.

Table 3: Types of Stomas and Associated Complications

Stoma Type	Frequency (n = 121)	Percentage (%)	Complication Rate (%)
Loop Stoma	71	58.7	25.4
End Stoma	14	11.6	21.4
Double-Barrel Stoma	36	29.8	38.9

This data supported the hypothesis that emergency procedures had more complications than elective procedures. Furthermore, patients with comorbid diseases with a focus on diabetes and ischemic heart disease had a higher risk of postoperative complications in the study. These observations underscore the need for proper management of patients before surgery and evaluation of these individuals after surgery, emphasizing high-risk groups.

DISCUSSION

To summarize, this study contributes by presenting the frequency of indications, and outcomes of enteric stoma formation in the surgical ward of Ayub Teaching Hospital Abbottabad. The results evidence of the demographic and clinical characteristics of the patients correspond to international trends in stoma delivery, especially in countries with high incidences of digestive pathologies and comorbidities resulting from trauma. Male dominance (65.3%) and the young mean age of 46.3 years indicate that stoma formation has economic significance since it happens in the working population. A wide range of indications for the creation of the stoma was identified in the present study, with intestinal obstruction being the most frequent reason observed in 36.4% of the cases. This is similar to the report of Mbelle et al. (2) in sub-Saharan Africa and Wahab et al. (7) in Malaysia, where delayed presentation and late stages of the disease often require stoma formation. Intestinal perforation, being the second most commonest indication (28.1%), was commonly seen in emergent cases, which was in concordance with the impoverished presentation indices among the local masses. These findings are in congruence with Sushel et al. (1), who showed that intestinal TB was a major cause of perforation in South Asia. External injuries that include abdominal trauma occurred in 20.7% of the patients and are indicative of the fact that external traumas are some of the conditions that call for emergency surgeries, which are in consonant with the findings of Chukwubuike et al. (11).

The type of stoma created depends on the disease and the type of emergency of the case. LOC was found in 58.7%, end stomas in 11.6% and double-barrel stomas in 29.8% of patients. These outcomes are quite passable to Granieri et al. (4), who also estimated similar great proportions in their systematic review. Loop stomas are preferred in temporary faecal diversion because of their simplicity in construction and reversal, especially in emergency situations. However, end stomas are usually made during the definitive surgery or when reversal will not be done, as observed by Nozawa et al. on stoma site marking and complication prevention. Issues arising from stomas remain a major challenge in the management of patients with stomas, as observed in this study, where 32.2% of the patients had postoperative complications. The minor complications encountered included wound infection, 15.7%, stoma retraction, 10.7%, and parastomal hernia, 5.8%. These rates are in concordance with Kimbugwe et al. (6), who also reported similar complications in their paediatric population, as well as Estrada et al. (9), who reported different layers of complications with Crohn's disease and ulcerative colitis. Data comparing complication rates between emergency and elective operations (42.3% vs 16.3%) support this claim, emphasizing better preoperative preparation and intraoperative and postoperative management of patients in emergencies.

Patients with comorbidity had a higher chance of complications with more risk posed to them if they had other diseases like diabetes or ischemic heart disease. For example, diabetic patients were more susceptible to wound infection and slow healing, as indicated in the work done by Tang et al. (8). This emphasizes the significance of glycemic management and close observation of the wounds in such patients. Likewise, patients with ischemic heart disease had higher mortality-morbidity risk after discharge after the surgery, thus the call for a comprehensive approach. Another important domain in the quality of stoma care is psychological health. Moya-Muñoz et al. (3) and Tang et al. (8) have previously reported high depression rates and lower quality of life among stoma patients. Although this study did not measure psychological outcomes, the results of this research have highlighted the future direction of psychological care embedding into stoma care.

The findings of this study also have implications for healthcare policy and resource allocation. This high proportion of emergencies, which represents 64.5% of all the cases, implies that people developed complications and sought medical attention only when they were significantly unwell, possibly because they could not access health care early or because they did not know they could. Investing in primary care, ensuring proper access to patients' first contact with the healthcare system and providing education to communities that address various early stages of gastrointestinal diseases minimizes the chance of needing emergency stoma creation. The issues which are identified based on the existing literature include failure to preoperatively mark the stoma site, as highlighted by Nozawa et al. (5). Adequate site marking can reduce problems such as retraction and skin irritation after surgery by a big margin, enhancing patient satisfaction. Adopting this practice as part of normal

conventional surgical procedures, including those presented in emergency situations, stands to benefit the patient immensely.

Other external factors have opportunities to improve treatment mechanisms, including the progress of surgical operations and patient post-surgery management. For example, according to Ahmed et al. (15), Simulation-based interventions can enhance the outcome of nurse performance and satisfaction with stoma care. The adoption of such training programs in local healthcare facilities can reduce disparities and make sure that patients get the best services. These results align with studies published in literature from other parts of the world done by Parini et al. (12) and Hapnes et al. (14). However, the localized data offer valuable information about the pertinent issues and concerns of healthcare providers in Pakistan, such as the high prevalence of intestinal tuberculosis, which underlines possibilities for focused intervention in these cases.

However, there are several limitations in this research that have to be acknowledged. There are certain drawbacks of non-propensity sampling, where quantitative samples are not representative of the population, and due to the small sample size, it is possible that some major problems related to stoma were not covered. Such studies in larger, multicenter trials are bound to give more discerning results and confirm these observations well. Finally, this study also provides a wide spectrum of indications for making enteric stoma, various types of stoma, and the results of enteric stoma formation in a tertiary care hospital in Pakistan. The results re-emphasize the global concept of early diagnosis, operative planning and post-operative management for better prognosis of the disease. Therefore, by addressing the above-outlined challenges and implementing best practices from different parts of the world, healthcare providers can improve the quality of stoma care, thereby minimizing the complications among the local population.

CONCLUSION

This paper aims to bring into focus the concept of enteric stoma for the management of gastrointestinal disorders with an insight into the various indicators, classifications, and results noticed in the surgical ward of Ayub Teaching Hospital Abbottabad. The most frequent indications of stoma formation were intestinal obstruction and perforation, with loop stomas being the most common. The majority of operations were carried out as emergencies; these cases are commonly related to a high incidence of complications. Diabetic patients and those with ischemic heart disease are some examples which showed a high-risk factor for postoperative complications, pointing to the fact that significant attention should be paid to preoperative and surgical site care. The study's results indicate a need for earlier diagnosis, improved patient enlightenment, and incisional planning, such as preoperative stoma site listing. Increased education of carers and the inclusion of psychology within stoma care practice is vital to fully meeting the patients' needs. Evaluation of these challenges is crucial in that healthcare systems will likely lessen the incidence of complications, enhance quality of life, and provide overall care for patients with stoma.

REFERENCES

- 1- Sushel, C., Abbas, K., Syed, B.M., Shaikh, S., Mallah, Q. and Pathan, M.R., 2022. Clinical Presentation and Surgical Management of Intestinal Tuberculosis Presented as an Acute Abdomen. *Journal of Liaquat University of Medical & Health Sciences*, 21(01), pp.1-5.
- 2- Mbelle, P.J., Kizigina, S. and Mwashambwa, M., 2024. Indications for Stoma Creation and Early Outcome Predictors of Stoma Closure at Universtiy of Dodoma Teaching Hospitals. *Int J Biomed Res Prac*, 4(2), pp.1-12.
- 3- Moya-Muñoz, N., Armenteros-Fernández, E., Bautista-Mártir, C., Vílchez-Díaz, I.D.P., López-Medina, I.M., Montoya-Juárez, R., Hueso-Montoro, C. and Capilla-Díaz, C., 2022. Assessment of health indicators in individuals with intestinal stoma using the nursing outcomes classification: A cross-sectional study. *Frontiers in Surgery*, 9, p.870379.
- 4- Granieri, S., Sessa, F., Bonomi, A., Paleino, S., Bruno, F., Chierici, A., Sciannamea, I.M., Germini, A., Campi, R., Talso, M. and Facciorusso, A., 2021. Indications and outcomes of

- enterovesical and colovesical fistulas: systematic review of the literature and meta-analysis of prevalence. *BMC surgery*, 21(1), p.265.
- 5- Nozawa, H., Sasaki, S., Hayashi, C., Kawasaki, A., Sasaki, K., Murono, K., Emoto, S. and Ishihara, S., 2024. Preoperative stoma site marking reduces postoperative stoma-related complications in emergency surgery: A single center retrospective cohort study. *Scandinavian Journal of Surgery*, 113(1), pp.40-49.
 - 6- Kimbugwe, G., Kakembo, N., Kisa, P., Okeny, P. and Sekabira, J., 2023. Prevalence of intestinal stoma complications and associated factors among children in Uganda, a cross-sectional study.
 - 7- Wahab, P.A., Mohamed, N., Ismail, N., Hassan, I.I. and Haryanto, H., 2024. A Descriptive Analysis of Patients with Stoma Attending a Tertiary Hospital on the East Coast of Peninsular Malaysia. *INTERNATIONAL JOURNAL OF CARE SCHOLARS*, 7(3), pp.4-10.
 - 8- Tang, W.S.W., Chiang, L.L.C., Kwang, K.W. and Zhang, M.W.B., 2022. Prevalence of depression and its potential contributing factors in patients with enterostomy: a meta-analytical review. *Frontiers in Psychiatry*, 13, p.1001232.
 - 9- Estrada, D.L., Magro, D.O., Benghi, L.M., Miranda, E.F., Ropelato, R.V. and Kotze, P.G., 2022. Postoperative Morbidity After Stoma Creation in Inflammatory Bowel Diseases: Differences Between Crohn's Disease and Ulcerative Colitis. *Journal of Surgery and Research*, 5(3), pp.468-474.
 - 10- Wuraola, F.O., Adesunkanmi, A.O., Mohammed, T.O. and Adisa, A.O., 2023. An Audit of Colostomy among Adult Patients in a Nigerian Tertiary Hospital. *Journal of Medical and Basic Scientific Research*, 4(3-4), pp.43-47.
 - 11- Chukwubuike, K.E., Okoloagu, N. and Eneh, W.U., 2023. Pediatric Stomas: A Study in a Teaching Hospital, Our Experience. *Journal of Clinical and Medical Reviews (JCMR)*, 1(1).
 - 12- Parini, D., Bondurri, A., Ferrara, F., Rizzo, G., Pata, F., Veltri, M., Forni, C., Coccolini, F., Biffi, W.L., Sartelli, M. and Kluger, Y., 2023. Surgical management of ostomy complications: a MISSTO–WSES mapping review. *World Journal of Emergency Surgery*, 18(1), p.48.
 - 13- Salah Abd Elaziz Soliman Elgendy, N., Shawky Mahamud, H., Mohamed Ahmed Ayed, M., Abo Elsoud Ahmed, H., Mohamed Samir, M. and Erian Shehata, M., 2025. Effect of Mind Mapping Application on Nurses' Performance regarding Colostomy at Pediatric Surgical Units. *Egyptian Journal of Health Care*, 16(1), pp.47-64.
 - 14- Hapnes, N.C., Stensvold, H.J., Bjørnland, K., Sæter, T., Guthe, H.J.T., Støen, R., Moltu, S.J., Rønnestad, A. and Klingenberg, C., 2024. Surgery for intestinal injuries in very preterm infants: a Norwegian population-based study with a new approach to disease classification. *BMJ Paediatrics Open*, 8(1), p.e002722.
 - 15- Ahmed Elbilgahy, A., Mohamed Ibrahim Nassar, H. and Ali Elsabely Mohammed, A., 2024. The Effect of Simulation-Based Intervention on Nurses' Performance and Satisfaction regarding Colostomy Care at Pediatric Surgical Unit. *Egyptian Journal of Health Care*, 15(1), pp.1450-1463.