



COMPARATIVE EFFECTS OF FARADIC AND RUSSIAN CURRENTS ON THE QUADRICEPS MUSCLE IN PATIENTS WITH KNEE OSTEOARTHRITIS

Kanika Bisht¹, Ojas Garg^{2*}, Kajal³, Anshul Singh Yadav⁴, Mahesh Kumar Paswan⁵

¹PG MPT Orthopedics, School of Medical and allied sciences, Sanskriti University, Uttar Pradesh 281401

²Assistant Professor, SCPM college of nursing and paramedical sciences, Atal Bihari Vajpayee Medical University, Uttar Pradesh 226002

³Assistant Professor, SCPM college of nursing and paramedical sciences, Atal Bihari Vajpayee Medical University, Uttar Pradesh 226002

⁴Assistant Professor, TS Mishra University, TS Mishra University, Uttar Pradesh – 226008

⁵Tutor, SCPM college of nursing and paramedical sciences, Atal Bihari Vajpayee Medical University, Uttar Pradesh 226002

***Corresponding Author:** Ojas Garg

*Assistant Professor, SCPM College of Nursing and Paramedical Sciences, Gonda (U.P).
Email – gargojus9@gmail.com

INTRODUCTION

Knee osteoarthritis (OA) joint disease commonly occurs in elderly population and it causes significant pain and functional limitations [1]. OA is classified into Primary OA and Secondary OA. Primary OA occurs as a result of natural wear and tear of knee joint structures due to aging, overuse or obesity, whereas Secondary OA occurs due to known primary causes like trauma, infection [2]. The Kellgren and Lawrence scale classifies OA into four grades as follows: Grade 0 indicates no radiographic findings of OA; Grade 1 indicates minimum osteophytes of doubtful clinical significance; Grade 2 indicates definitive osteophytes with unaffected joint space; Grade 3 indicates definitive osteophytes with moderate joint space narrowing; and Grade 4 indicates definitive osteophytes with severe joint space narrowing and subchondral sclerosis [3]. Pathologically it is characterized by focal loss of articular hyaline cartilage with proliferation of new bone at the margins, subchondral sclerosis and remodeling of joint contour. The earliest symptom of OA knee is pain. Patients also complain of crepitus, stiffness, swelling of the joint. On examination, tenderness along the joint line, terminal limitation of movement, weakness and atrophy of the quadriceps femoris muscle. The radiographic features of OA knee include narrowing of joint space, osteophyte formation, loose bodies and subchondral sclerosis [5]. Joint space narrowing occurs due to the erosion of the cartilage as part of the degenerative process associated with arthritis but the best method of evaluating the progression of the cartilage destruction is through measurement of joint space width [6].

A recent survey in India reported that the prevalence of OA in older adults more than 65 years of age was 32.6% in the pastoral population and 60.3% in the urban population [7]. Men are affected more commonly than women [8].

Various measures are available for the treatment of OA knee such as conservative management includes pharmacotherapy (NSAIDs, Corticosteroid therapy), physiotherapy and surgical

management [5]. Manual therapy, physical therapy modalities, taping techniques, patient education and therapeutic exercises, and orthosis are all used to treat knee OA [9].

According to the World Journal of Orthopedics 2011, isometric exercises are helpful in improving muscle function. Exercises such as static Quadriceps, static hamstrings, strengthen the vastus medialis, mini squats, use of Quadriceps table, 10 to 15 repetitions, done 5 days a week for 2 weeks have proved to be effective in improving muscle strength [10,11]. Quadriceps femoris muscle plays an important role in mechanics of knee joint [12]. It is the only muscle crossing anterior to the axis of the knee and is the prime mover for knee extension. During quiet standing and stance phase of gait, knee is an intermediate joint in a closed chain. The quadriceps muscle controls the amount of flexion at the knee during initial contact and also causes knee extension through reverse muscle pull on the femur during mid-stance. It also controls the amount of flexion during pre-swing (heel off to toe off) and excessive heel rise during initial swing. With loss of quadriceps function, the patient lurches the trunk anteriorly during initial contact to move the center of gravity anterior to the knee so that it is stable or rotates the extremity outward to lock the knee; with fast walking there may be extreme heel elevation during initial swing [13].

Russian current is a medium frequency current. Russian current was developed for upgrading muscle strength in Russian Olympic athletes and was found to increase force up to 40%. Russian current is an Alternating Current modulated sinusoidal with 2500-Hz transfer in a series of bursts (10-millisecond burst and a 10-millisecond interval). It has been reported that the most commonly used electric stimulation to increase muscle strength, reduce pain and improve physical function is Russian current [14].

Neuromuscular stimulation may be an alternative approach for muscle strengthening training program. Faradic current may be the better mode of electrical stimulation to boost muscle strength and muscle mass. Voluntary muscle contraction followed by faradic current may result in functional faradism at the same time. Functional faradism is an essential portion of neuromuscular stimulation which can be useful in boosting muscle mass, strength of muscle and improve functional mobility in knee OA. To improve the muscle strength and size of muscle fiber, each contraction of muscle followed by faradic stimulation simultaneously may be better and useful rehabilitative way of knee OA. Although previous studies found improvement in muscle mass, strength of muscle and function after implementation of faradic current stimulation [15]. Surged faradic current is used for the upgrading muscle strength and vascularity of muscle [16].

Purpose of the study:

Many Previous studies investigated the effectiveness of Russian current and also faradic current on the quadriceps muscle in patients with knee osteoarthritis but no any comparative study was performed between the effect of Faradic and Russian current on the quadriceps muscle for management of knee osteoarthritis. So, this study will conduct to examine the Comparison between effect of Faradic and Russian current on the quadriceps muscle for management of knee osteoarthritis.

Significance of the study:

This study will be conducted to find out the comparative effect of Russian and faradic current for the management of OA knee and this study will be beneficial to create better physiotherapy treatment protocols for the management of OA knee and this study aims to provide important information about the management of OA and to impart knowledge.

Aims of the study:

- To find out the Comparison between Faradic and Russian Currents on the quadriceps muscle in patients with knee osteoarthritis.

Objectives:

- To evaluate the effect of Faradic Currents on the quadriceps muscle for management of knee osteoarthritis.
- To evaluate the effects of Russian Currents on the quadriceps muscle for management of knee osteoarthritis.
- Comparative effects between Faradic and Russian Currents on the quadriceps muscle for management of knee osteoarthritis.

Research hypothesis

- **Null hypothesis (H_0):** - There will be no significant difference between Faradic and Russian current on the quadriceps muscle in patients with knee osteoarthritis.
- **Experimental hypothesis (H_1):** - There will be significant difference between Faradic and Russian current on the quadriceps muscle in patients with knee osteoarthritis.

METHODOLOGY**Sample size:**

A sample of 30 subjects (2 groups and 15 subjects in each group) was selected to take part in the study based on the fulfillment of inclusion and exclusion criteria.

Inclusion criteria:

- Pre-Diagnosed case of Grade II OA patients
- Age: 45-60yr
- Gender: Both male & female
- Complaining of knee pain that has lasted longer than 3 months,
- A pain level that is medium

Exclusion criteria:

- Any recent hip or knee surgeries & fractures Patients with neurological deficit
- Soft tissue injuries of lower limb
- Any systemic illness
- Red flag such as trauma, cancer, TB spine

Study Design:

A Randomized Clinical Trial.

VARIABLES**Dependent:**

- 1) NPRS (Numeric pain rating scale),
- 2) Goniometric measurement for ROM of knee,
- 3) Timed Up and Go (TUG) test,
- 4) Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC)

Independent:

- 1) Faradic current
- 2) Russian current

Instrumentation:

- 1) A stopwatch
- 2) Standard Chair with arm rest
- 3) Universal Goniometer
- 4) Weighing machine
- 5) A stadiometer

- 6) Couch
- 7) Electrical stimulation machine having Faradic Current
- 8) Electrical stimulation machine having Russian Current

Outcome measures:


1) **NPRS (Numeric pain rating scale):** It is a unidimensional measurement tool use for measure the pain intensity in adults. The NPRS is a segmented numeric version of the visual analog scale (VAS) in which a respondent selects a whole number (0-10 integers) that best display the pain intensity of the patient. The common format is a horizontal line and 11- point numeric scale ranges from _0'(no pain) to _10'(Worst pain imaginable) [18, 19].


Figure 3.1: Numeric pain rating scale (NPRS)

2) **Goniometry of knee ROM:** Knee ROM is commonly used as an outcome measure in clinical trials of people with knee OA. Goniometry is a reliable and valid way to measure knee ROM, and is a convenient, accessible outcome measure for clinical trials and physiotherapy practice [33]. Knee flexion and extension occurs in the sagittal plane around a medial-lateral axis. The normal range of flexion for adults varies from 130 to 140 degrees and the normal knee extension ROM values for adults are about 0 degrees. When extension goes beyond the 0-starting position, it may be within the normal limits of 5 to 10 degrees in children, but when it exceeds 5 or more degrees in adults, it is called hyperextension [34].

3) **Timed Up and Go (TUG) test:** The TUG test is one of the simple and quick tests to assess patients' functional mobility test that was originally developed for frail, elderly people, but is now also recommended for musculoskeletal conditions, such as hip and knee osteoarthritis (OA) The TUG is a transition test that assesses strength, agility and dynamic balance during multiple activities including sit-to-stand, walking short distances and changing direction during walking. Most healthy people upto the age of 80 are able to perform the TUG in 10 seconds or less [35, 36].

4) **Western Ontario and McMaster Universities Osteoarthritis Index:** The WOMAC Osteoarthritis Index is a disease-specific self-report questionnaire for measurement of the symptoms of OA of the hips and knees. It is reliable, valid, and sensitive to the changes in the health status of patients with knee OA. We used the 3.1 Likert version with five response levels for each item, representing different degrees of intensity (none, mild, moderate, severe, or extreme) that were scored from 0 to 4. The final score for the WOMAC was determined by adding the aggregate scores for pain, stiffness, and function. Scores range from 0 to 96 for the total WOMAC where 0 represents the best health status and 96 the worst possible status. The higher the score, the poorer the function [37].





PROTOCOL: Figure 3.2: Consort flow chart of study

PROCEDURE

A total of 40 subjects were enrolled in this study. Both male and female were participated in this study having 45 -60 years. The subjects made to sign the inform consent form before the study. The consent form was explained in the language which comfortable for the patient. 30 subjects were randomly selected in the treatment groups (Group A&B) by simple random sampling by chit picking method on the basis of inclusion and exclusion criteria. Participant's demographic data was collected including their bodyweight in kg measured by using the weighing machine, Height in cm measured by using the stadiometer and Participant's BMI which is in Kg/m² was calculated using the participant's weight and dividing it by the square of the appropriate height. Baseline measurements for pain intensity by NPRS, knee ROM by goniometer, Timed Up and Go (TUG) test and WOMAC were taken for every patient after that faradic currant + Quadriceps isometric was applied for group A and Russian current + Quadriceps isometric was applied for group B than post intervention outcome measures were taken at the end of 4th week.




Figure 3.3: Height Measured by Stadiometer.

Baseline and post intervention measurements procedure:

NPRS (Numeric pain rating scale): Baseline and after 4th week measurements for knee pain intensity by NPRS in which asked the patients to select the numerical value of the scale i.e., range 0-10 and express verbally which indicates about his/her pain intensity [18].

Range of Motion of Knee: Baseline and after 4th week measurements for active range of motion of knee was measured by a universal goniometer and recorded in degree. These measurements were taken for knee flexion and knee extension. According to Norkin & white when measuring the knee flexion, the patient placed in supine lying with the knee extension. Position the hip in 0 degrees of extension, abduction and adduction. The femur Stabilization was done to prevent rotation, abduction, and adduction of the hip. The testing position was hold the anterior thigh with one hand and ankle with the other hand Move the patient's thigh to approximately 90 degrees of hip flexion and move the knee into flexion. To prevent further hip motion, stabilize the thigh and guide the lower leg into knee flexion. When resistance is felt the end of the range of knee flexion was occurs. Goniometer placement was, fulcrum placed over the lateral epicondyle of the femur, Stationary arm with the lateral midline of the femur, using greater trochanter as a reference and movement arm with the lateral midline of the fibula, using the lateral malleolus and fibular head as a reference and during measurement of knee extension the patient placed in supine lying. A folded towel placed under the ankle to ensure that the knee is in full extension. The femur Stabilization was done to prevent rotation, abduction, and adduction of the hip. Goniometer placement in knee extension measurement is same as knee flexion [34].



Figure 3.4: Goniometry for Knee ROM.



Figure 3.5: Timed Up and Go (TUG) test.

Timed Up and Go (TUG) test: Baseline and after 4th week measurements for TUG test subjects were instructed to stand up from the chair, walk 3 meters comfortably and safely than come back and sit in the chair. The time taken to complete this task was measured with a stopwatch [36].

WOMAC Arthritis Index: The Western Ontario and McMaster Universities Arthritis Index was done before (pre) and after (post) the intervention, including five response levels for each item, representing different degrees of intensity (none, mild, moderate, severe, or extreme) that was scored from 0 to 4. The pain dimension or scale includes five items asking pain at activity or rest. The stiffness dimension includes two questions. The function dimension explores the degree of difficulty in 17 activities. The final score for the WOMAC was determined by adding the aggregate scores for pain, stiffness, and function. Scores range from 0 to 96 for the total WOMAC where 0 represents the best health status and 96 the worst possible status. The higher the score, the poorer the function. Therefore, an improvement was achieved by reducing the overall score [37]

Intervention:

Group A: Faradic current + quadriceps isometric exercise (5days in a week for 4 week)

After taken the baseline measurement of the patient we started the intervention of the patients who are randomly selected in group A than applied the faradic current after the resting period of 10 min following quadriceps isometric exercise. For faradic stimulation the patient were placed comfortable in supine lying position on a couch with pillow placed below the knee joint adequate positioning of standard carbon rubber electrodes of equal paired size were placed on the quadriceps muscle of the arthritic knee joint .the electrodes was securely fixed using Velcro strap and set the electrical stimulation intensity which was set at the maximum tolerance of the patient. Parameters included rectangular biphasic symmetric current, pulse duration of 400 μ s and stimulation frequency of 80 Hz. The session was given for 10 min and was comprised of 10 s of stimulation followed by 20 s of rest for 5 days/week for 4 weeks. For quadriceps isometric exercise Patients were in supine position with legs straight with the towel roll placed under the knee, patients were asked to press the knee on the towel roll. This exercise aimed to strengthen quadriceps muscle. The patient was instructed to maximally activate their thigh muscles in order to straighten their knee. This exercise was of 3 sets of 10 repetitions each for 5 days/week for 4 weeks (20 days).



Figure 3.6: Faradic current stimulation.

Group B: Russian current + quadriceps isometric exercise (5 days in a week for 4 weeks).

After taken the baseline measurement of the patient we started the intervention of the patients who are randomly selected in group B than applied the Russian current after the resting period of 10 min following quadriceps isometric exercise. For Russian stimulation the patient were placed comfortable in supine lying position on a couch with pillow placed below the knee joint adequate positioning of standard carbon rubber electrodes of equal paired size were placed on the quadriceps muscle of the arthritic knee joint. the electrodes was securely fixed using Velcro strap and set the electrical stimulation intensity which was set at the maximum tolerance of the patient. Russian Current of 2500 Hz with pulse frequency of 50 Hz, symmetric pulses of sinusoidal form, with a pulse duration of 200-300 μ s, 2-3 seconds ramp up, 2 seconds ramp down for comfort and a duty cycle of 50% was applied for 10 minutes duration for 5 days/week for a period of 4 weeks after that instructed the patient to quadriceps isometric exercise. This exercise was of 3 sets of 10 repetitions each for 5 days/week for 4 weeks.



Figure 3.7: Russian current stimulation.

DATA ANALYSIS

Analysis was done for 30 subjects who completed the study. The outcome variables of the study included Knee pain intensity on NPRS, Knee range of motion by Goniometry, TUG and WOMAC. Paired t-test was used for comparing the pretreatment and post-treatment scores of each variable for both the groups (within group analysis). Independent t-test was applied to compare the outcome values between the group A and group B. The value of all two groups i.e., Group A (Faradic current) and Group B (Russian current), were compared at the pretest and after 4th Week of treatment.

Statistical significance was set at $P < 0.05$. P value > 0.05 was considered as non-significant difference while P value < 0.05 was considered to have represented a significant difference. Value of confidence interval was set at 95%.

RESULT

The statistics done based on the objectives and hypotheses and it has been organized by following three sections.

Section I: Frequency and percentage distributions of sociodemographic variables.

Section II: Effectiveness of pre and posttest values.

Section III: Comparison of Faradic Currents and Russian Currents values.

Section I: Frequency and percentage distributions of sociodemographic variables.

Table 5.1: Frequency and percentage distributions of sex

Sociodemographic Data	n=15		n=15	
	Faradic Currents	Russian Currents	Faradic Currents	Russian Currents
	Frequency	Percentage	Frequency	Percentage
Sex				
1) Male	7	46.7	7	46.7
2) Female	8	53.3	8	53.3

The above table implies that maximum 8(53.3%) samples in faradic currents as well as in Russian currents are females, 7(46.7%) of the samples are males in each.




Figure 5.1: Frequency and percentage distributions of sex

Table 5.2: Frequency and percentage distributions of age

Sociodemographic Data	n=15		n=15	
	Faradic Currents	Russian Currents	Faradic Currents	Russian Currents
	Frequency	Percentage	Frequency	Percentage
Age in years				
1) 45 to 49	7	46.7	5	33.3
2) 50 to 54	4	26.7	5	33.3
3) 55 to 60	4	26.7	5	33.3

The above table implies that in faradic currents the maximum 7(46.7%) samples are between the age 45 to 49 years, 4 (26.7%) samples are between 50 to 54 years and the remaining 4(26.7%) samples are between 55 to 60 years. Whereas in Russian currents 5(33.3%) samples are in between 45 to 49 years, another 5(33.3%) samples are in 50 to 54 years and the remaining 5(33.3%) samples are between 55 to 60 years.




Figure 5.2: Frequency and percentage distributions of age

Table 5.3: Frequency and percentage distributions of Weight (kg)
n=15 n=15

Sociodemographic Data	Faradic Currents		Russian Currents	
	Frequency	Percentage	Frequency	Percentage
Weight (Kg)				
1) 51 to 59	1	6.7	2	13.3
2) 60 to 68	8	53.3	3	20.0
3) 69 to 77	5	33.3	3	20.0
4) 78 to 86	1	6.7	5	33.3
5) 87 to 95	0	0	1	6.7
6) 96 to 104	0	0	1	6.7

The above table explains that in the faradic currents the maximum 8 (%53.3) samples weight was between 60 to 68 kg, whereas in Russian currents the maximum 5 (33.3%) samples weight is between 78 to 86 kg.





Figure 5.3: Frequency and percentage distributions of Weight (kg)

Table 5.4: Frequency and percentage distributions of Height (cm)
n=15 n=15

Sociodemographic Data	Faradic Currants		Russian Currants	
	Frequency	Percentage	Frequency	Percentage
Height				
1) 150 to 155	2	13.3	4	26.7
2) 156 to 161	4	26.7	2	13.3
3) 162 to 167	6	40.0	4	26.7
4) 168 to 173	3	20.0	5	33.3

The above table implies that in the faradic currants the maximum 6(40.0%) samples height is between 162 to 167, whereas in Russian currants maximum 5(33.3%) samples height is between 168 to 173.

**Figure 5.4: Frequency and percentage distributions of Height (cm)****Table 5.5: Frequency and percentage distributions of BMI (kg/m²)**
n=15 n=15

Sociodemographic Data	Faradic Currants		Russian Currants	
	Frequency	Percentage	Frequency	Percentage
BMI (kg/m ²)				
1) 22 to 27	13	86.7	6	40.0
2) 27 to 32	2	13.3	7	46.7
3) 32 to 37	0	0	2	13.3

The above table implies that in the faradic currants the most 13(86.7%) samples BMI (kg/m²) is 22 to 27 whereas in Russian currants the maximum 7(46.7%) samples BMI (kg/m²) is 27 to 32.




Figure 5.5: Frequency and percentage distributions of BMI (kg/m²)

Section II: Effectiveness of pre and posttest values

Table 5.6: Effectiveness of pre and posttest Faradic Currants values

n=15					
Pre-test and Post-test Faradic Aspects	Standard Error Mean	Mean	SD	Students' paired t-test	Significance
Pre Pain - Post Pain	0.192	3.133	0.743	16.328	0.000; P<0.05; S
Pre Knee-flexion - Post Knee flexion	1.059	7.867	4.103	7.425	0.000; P<0.05; S
Pre Knee-Extension - Post Knee Extension	.452	9.933	1.751	21.969	0.000; P<0.05; S
Pre-TUG - Post TUG	.279	3.200	1.082	11.451	0.000; P<0.05; S
Pre WOMAC - Post WOMAC	.487	7.533	1.885	15.480	0.000; P<0.05; S

S=Significant; SD= Standard deviation; df=14

The above Table depicts the calculated 't' value of each aspects are higher than the tabulated value at 0.05 level of significance. So, there is a significant difference between pretest and posttest Faradic Currants values.

Table 5.7: Effectiveness of pre and posttest Russian Currants values

n=15					
Pre-test and Post-test Russian Aspects	Standard Error Mean	Mean	SD	Students' paired t-test	Significance
Pre Pain - Post Pain	0.133	3.533	.516	26.500	0.000; P<0.05; S
Pre Knee-flexion - Post Knee flexion	1.018	7.467	3.944	7.333	0.000; P<0.05; S
Pre Knee-Extension - Post Knee Extension	1.018	10.867	3.944	10.672	0.000;

					P<0.05; S
Pre-TUG - Post TUG	.223	4.200	.862	18.873	0.000; P<0.05; S
Pre WOMAC - Post WOMAC	.696	10.400	2.694	14.952	0.000; P<0.05; S

S=Significant; SD= Standard deviation; df=14

The above Table depicts the calculated 't' value of each aspects are higher than the tabulated value 2.145 at 0.05 level of significance. So, there is a significant difference between pretest and posttest Russian Currants values.

Section III: Comparison of Faradic Currants and Russian Currants values

Table 5.8: Effectiveness of Faradic Currants and Russian Currants in Pre and Post Pain values

n=15+15					
Faradic Currants and Russia Currants Aspects	Standard Error Mean	Mean	SD	Independent t-test	Significance
Pre Pain Faradic Currants	.192	6.47	.743	0.498	P>0.05; NS
Pre Pain Russian Currants	.187	6.33	.724		
Post Pain Faradic Currants	.232	3.33	.900	1.740	P>0.05; NS
Post Pain Russian Currants	.200	2.80	.775		

n=15+15						
Faradic Currants and Russian Currants Aspects	Standard Error Mean	Mean	SD	Independent t-test	Significance	
Pre Knee Extension Faradic Currants	1.080	10.93	4.183	0.228	P>0.05; NS	
Pre Knee Extension Russian Currants	1.380	10.53	5.343			
Post Knee Extension Faradic Currants	1.000	1.00	3.873	0.768	P>0.05; NS	
Post Knee Extension Russian Currants	1.420	.33	5.499			

The above table depicts that the t test value of pre pain and post pain in Faradic currants and Russian currants are lower than the tabulated value 2.048 at 0.05 level of significance. So, there is no significant difference between Faradic Currants and Russian Currants in Pre and post Pain values.




Figure 5.6: Effectiveness of Faradic Currants and Russian Currants in Pre and Post Pain values

Table 5.9: Effectiveness of Faradic Currants and Russian Currants in pre and post Knee flexion values

						n=15+15
Faradic Currants and Russian Currants Aspects	Standard Error Mean	Mean	SD	Independent t-test	Significance	
Pre Knee flexion Faradic Currants	1.291	119.47	4.998	0.469	P>0.05; NS	
Pre Knee flexion Russian Currants	1.515	118.53	5.866			
Post Knee flexion Faradic Currants	.826	127.33	3.200	0.962	P>0.05; NS	
Post Knee flexion Russian Currants	1.113	126.00	4.309			

The above table depicts that the t test value of pre and post Knee flexion in Faradic currants and Russian currants are lower than the tabulated value 2.048 at 0.05 level of significance. So, there is no significant difference between Faradic Currants and Russian Currants in pre and post Knee flexion values.





Figure 5.7: Effectiveness of Faradic Currants and Russian Currants in pre and post Knee flexion values

Table 5.10: Effectiveness of Faradic Currants and Russian Currants in pre and post Knee Extension values

The above table depicts that the t test value of pre and post Knee Extension in Faradic currants and Russian currants are lower than the tabulated value 2.048 at 0.05 level of significance. So, there is no significant difference between Faradic Currants and Russian Currants in pre and post Knee Extension values.

**Figure 5.8: Effectiveness of Faradic Currants and Russian Currants in pre and post Knee flexion values****Table 5.11: Effectiveness of Faradic Currants and Russian Currants in pre and post TUG values**

n=15+15					
Faradic Currant&Russian Currants Aspects	Standard Error Mean	Mean	SD	Independent t-test	Significance
Pre TUG Faradic Currants	.262	16.20	1.014	0.458	P>0.05; NS
Pre TUG Russian Currants	.349	16.40	1.352		
Post TUG Faradic Currants	.309	13.00	1.195	1.871	P>0.05; NS
Post TUG Russian Currants	.296	12.20	1.146		

The above table depicts that the t test value of pre and post TUG in Faradic currants and Russian currants are lower than the tabulated value 2.048 at 0.05 level of significance. So, there is no significant difference between Faradic Currants and Russian Currants in pre and post TUG values.




Figure 5.9: Effectiveness of Faradic Currants and Russian Currants in pre and post TUG values

Table 5.12: Effectiveness of Faradic Currants and Russian Currants in pre and post WOMAC values

n=15+15					
Faradic Currants and Russian Currants Aspects	Standard Error Mean	Mean	SD	Independent t-test	Significance
Pre WOMAC Faradic Currants	.947	41.80	3.668	1.370	P>0.05; NS
Pre WOMAC Russian Currants	.910	40.00	3.525		
Post WOMAC Faradic Currants	1.201	34.27	4.652	3.172	P<0.05; S
Post WOMAC Russian Currants	.850	29.60	3.291		

S=Significant; NS=Not Significant SD= Standard deviation; df=28

The above Table depicts the calculated 't' value of Pre WOMAC Faradic and Russian Currant t test values are lower than the tabulated value 2.048 at 0.05 level of significance. So, there is no significant difference between Faradic Currants and Russian Currants pre WOMAC values. In the Post WOMAC Faradic Currants and Post WOMAC Russian Currant t test values are higher than the tabulated value 2.048 at 0.05 level of significance. So, there is a significant difference between Faradic Currants and Russian Currants WOMAC values.




Figure 5.10: Effectiveness of Faradic Currants and Russian Currants in pre and post WOMAC values

DISCUSSION

The study was designed to determine the effect of Faradic and Russian current on the quadriceps muscle in patient with knee OA. The purpose of the study was to compare the effect of Faradic and Russian current on quadriceps muscle in patient with knee OA in order to increase ROM of the knee, decrease pain and improve functional ability of knee joint. The findings of this study are based on the objectives and hypotheses and it has been organized by following three sections.

Section I: Frequency and percentage distributions of sociodemographic variables.

The present study shows that the maximum 8(53.3%) samples in faradic currants as well as in Russian currants are females, 7(46.7%) of the samples are males in each. In faradic currants the maximum 7(46.7%) samples are between the age 45 to 49 years, 4 (26.7%) samples are between 50 to 54 years and the remaining 4(26.7%) samples are between 55 to 60 years. Whereas in Russian currants 5(33.3%) samples are in between 45 to 49 years, another 5(33.3%) samples are in 50 to 54 years and the remaining 5(33.3%) samples are between 55 to 60 years. In the faradic currants the maximum 8 (%53.3) samples weight was between 60 to 68 kg, whereas in Russian currants the maximum 5 (33.3%) samples weight is between 78 to 86 kg. In the faradic currants the maximum 6(40.0%) samples height is between 162 to 167, whereas in Russian currants maximum 5(33.3%) samples height is between 168 to 173. In the faradic currants the most 13(86.7%) samples BMI (kg/m²) is 22 to 27 whereas in Russian currants the maximum 7(46.7%) samples BMI (kg/m²) is 27 to 32.

Section II: Effectiveness of pre and posttest values.

The present study revealed the there is a significant difference between pretest and posttest Faradic Currants values and also there is a significant difference between pretest and posttest Russian Currants values. It means Faradic and Russian current both are effective on quadriceps muscle in patient with knee osteoarthritis in order to increase ROM of the knee, decrease pain and improve functional ability of knee joint.

Section III: Comparison of Faradic Currants and Russian Currants values.

In this study we found that there is no significant difference between Faradic Currants and Russian Currants in Pre and post Pain (NPRS) values, in pre and post Knee flexion values, in pre and post Knee Extension values, in pre and post TUG values and in pre WOMAC values (Therefore, the H_0 is accepted and the H_1 is rejected) but there is a significant difference between Faradic Currants and Russian Currants post WOMAC values (Therefore, the H_0 is rejected and the H_1 is accepted).

A study conducted by Vaz et al on neuromuscular electrical stimulation (NMES) reduces structural and functional losses of quadriceps muscle and improves health status patients with knee osteoarthritis. This study concluded that the NMES strength training program was successful in

counteracting the deleterious effects of OA on the knee extensor muscles. NMES reduced muscle weakness and increased muscle thickness and fascicle length. NMES training appears to offset the changes in quadriceps structure and function, as well as improve the health status in patients with knee OA [15].

The previous study conducted by dr. raghav s. et al to determine the comparative effect of the functional faradism over strengthening exercises in reducing pain, joint stiffness, functional limitation and improving isometric quadriceps muscle strength and quality of life in patients with knee osteoarthritis. This study revealed that the reduction of pain, joint stiffness and functional limitation in both groups may be attributed to increased quadriceps muscle strength and thereby improve function which leads to decrease pain and disability. The results of this study shown that the functional faradism was an effective protocol for reducing pain, joint stiffness, functional limitation and improving quadriceps muscle strength [25].

According to Ward AR et al the effect of Russian current stimulation as it is believed to relieve pain, re-educates muscle function, prevents muscle atrophy and restores function. Russian current can cause depolarization of sensory and motor nerve fibers. It activates fast type II motor unit and evoke muscle contraction which leads strengthening of muscle. [14]

Michael G Parker et al. conducted a study on torque responses in human quadriceps to burst modulated alternating current at 3 carrier frequencies namely, 2500 Hz, 3750 and 5000 Hz in 23 healthy subjects, concluded that burst modulated alternating current given at 2500 Hz produces greater electrically induced torque than those generated at 3750 Hz and 5000 Hz. According to this study we used Russian current stimulator having 2500 Hz frequency in present study [38].

According to previous study the improvement in function could be attributed to analgesic effect of Russian current and exercises which lead to decrease pain and increase strength of quadriceps. Strength gain was also associated with improvement in confidence during mobility [39].

Some previous study revealed that Russian current stimulation of the peripheral muscle causes an increase the spinal routes excitability, changes in the cortical activation pattern and improves the recruitment of fast twitch (fatigable) muscle fibers which are responsible for strength [40]. Also, there could be some neural adaptations that increase the capacity of voluntary muscle contraction that was impaired in patients with knee OA [41].

A study conducted by Anand B.H, Snehal R.D et al on effect of Russian current on questions Quadriceps muscle strength with primary knee osteoarthritis. This study concluded that the Russian current group Shows better improvement in pain and increase in muscle strength and greater improvement of the functional score (WOMAC) [26].

LIMITATIONS:

There are several limitations to this study. The sample size in present study was small, No follow up was done, strength of the quadriceps muscle was not included, and Radio graphical findings were not measured in this study.

RECOMMENDATIONS FOR FUTURE STUDY:

Further studies may be done with larger sample size to generalize the results. Long term follow up of the patients is recommended in further studies to see the long term effects. Radiographical findings are recommended in further studies to measure the effects.

CONCLUSION

This study conclude that faradic current and Russian current show significant effect in quadrics muscle in patients with knee osteoarthritis and there is no significant difference between faradic and Russian current on quadriceps muscle Patients with knee osteoarthritis. But only significant difference in WOMAC scores after intervention (The WOMAC score is more decrease in Russian as compare to faradic current)

BIBLIOGRAPGY

1. **Bricca A, Juhl CB, Steultjens M et al (2019)** Impact of exercise on articular cartilage in people at risk of, or with established, knee osteoarthritis: a systematic review of randomised controlled trials. *Br J Sports Med* 53(15):940–947. <https://doi.org/10.1136/bjsports-2017-098661>.
2. **Jayant Joshi, Prakash Kotwal.** Essentials of Orthopedics and Applied Physiotherapy. Elsevier Pub. 2011: 2nd edition, Chapter.16:341.
3. **Kellgren, J. H.; Lawrence, J. S. (1957).** Radiological Assessment of Osteo-Arthrosis. *Annals of the Rheumatic Diseases*, 16(4), 494–502.
4. **Boon et al** Davidson principle and practice of medicine, 20th edition. 2006(13): 1096-1097.
5. **J. Maheshwari.** Essential Orthopedics. Jaypee Brothers 2011: 4th edition Chapter. 35: 287-289.
6. **E. Vignon, M. Piperno, M.P. Hellio Le Graverand, S.A. Mazzuca, et al** “Measurement of Radiographic Joint Space Width in the Tibiofemoral Compartment of the Osteoarthritic Knee. Comparison of standing anteroposterior and Lyon Schuss views.” *Arthritis and Rheumatism*, 48(2): 378-384, 2003.
7. **Sharma MK, Swami HM, Bhatia V, et al.** An epidemiological study of correlates osteoarthritis in geriatric population of UT Chandigarh. *Indian J Community Med*, 2007, 32: 77–78.
8. **Robin McKenzie, Stephen May.** The human extremities –Mechanical Diagnosis and therapy. Spinal Publication New Zealand Ltd 2007 1st edition: 255 -260.
9. **de Oliveira MM, Aragao FA, Vaz MA (2013)** Neuromuscular electrical stimulation for muscle strengthening in elderly with knee osteoarthritis - a systematic review. *Complement Ther Clin Pract* 19(1):27–31.
10. **Jun Iwamoto et al.** Effectiveness of exercise for osteoarthritis of the knee: A review literature. *World Journal of Orthopedics*, 2011 May 18; 2(5):37-42.
11. **Rene Calliet.** Knee pain and disability. Jaypee Brothers; 3rd edition 190-197.
12. **Michael V. Hurley.** The role of muscle weakness in the pathogenesis of osteoarthritis. *Rheumatic Disease Clinics of North America*, 1 May 1999; 25(2): 283-298.
13. **Carolyn Kisner, Lynn Allen Colby.** Therapeutic Exercise Foundations and Techniques. Jaypee Brothers 2007: 5th edition. Chapter 21: 691-693.
14. **Ward AR, Shkuratova N.** Russian electrical stimulation: the early experiments. *Physical therapy*. 2002 Oct 1; 82(10):1019-30.
15. **Marco Aurélio Vaz et al. (2012),** Neuromuscular electrical stimulation (NMES) reduces structural and functional losses of quadriceps muscle and improves health status in patients with knee osteoarthritis. *Journal of Orthopaedic Research* Volume 31, Issue 4: 511-516.
16. **Ratan SK, Rattan KN (2009).** The surged faradic stimulation to the pelvic floor muscle as an adjunct to the medical management in children with rectal prolapsed, *BMC pediatrics*.
17. **International Association for the Study of Pain (2011).** Classification of Chronic Pain. *Part III Pain Therms.* 2.
18. **Jensen, M., & McFarland, C. (1993).** Increasing the reliability and validity of pain intensity measurement in chronic pain patients. *Pain*, 55(2), 195-203.
19. **Childs, J., Piva, S., & Fritz, J. (2005).** Responsiveness of the Numeric Pain Rating Scale in Patients with Low Back Pain. *Spine*, 30(11), 1331-1334.

20. **S. Jagmohan (2018)**, Textbook of Electrotherapy, Jaypee Brothers Medical Publishers (P) Ltd, 3rd edition, Chapter 2:75.
21. **S. Jagmohan (2018)**, Textbook of Electrotherapy, Jaypee Brothers Medical Publishers (P) Ltd, 3rd edition, Chapter 3:134.
22. **Norkin C. C., D. J. White (2016)**, Measurement of joint motion: a guide to goniometry, Philadelphia : F.A. Davis Company, Fifth edition, Chapter 1:7
23. **John O. Omole, Michael O. Egwu, et al (2018)**, Comparative Effects Of Russian Current And Isometric Resisted Exercise On Quadriceps Angle And Joint Space Width Among Patients With Primary Knee Osteoarthritis, *International Journal of Advanced Research and Publications*, Vol 2(2): 92-99.
24. **S. Mathew et al (2020)**. Effectiveness of eccentric exercise and neuromuscular electrical stimulation on quadriceps function in grade II osteoarthritis- a randomized control trial, *International Journal of Research and Review* Vol.7; Issue: 4; 328-335
25. **Dr. Raghav S, Dr. A Singh (2020)**, comparative effect of the functional faradism over strengthening exercises in reducing pain, joint stiffness, functional limitation and improving isometric quadriceps muscle strength and quality of life in patients with knee osteoarthritis, *International Journal of Creative Research Thoughts (IJCRT)*, 8(4); 2948-2956.
26. **Anand B Heggannavar, Snehal R Dharmayat, Sonal S Nerurkar (2014)**, EFFECT OF RUSSIAN CURRENT ON QUADRICEPS MUSCLE STRENGTH IN SUBJECTS WITH PRIMARY OSTEOARTHRITIS OF KNEE: A RANDOMIZED CONTROL TRIAL. *International Journal of Physiotherapy and Research, Int J Physiother Res*, Vol 2(3):555-60.
27. **Rosemffet, Marcos G.; Schneeberger, Emilce E.; Citera, Gustavo, et al. (2004)**. Effects of Functional Electrostimulation on Pain, Muscular Strength, and Functional Capacity in Patients With Osteoarthritis of the Knee. *JCR: Journal of Clinical Rheumatology*, 10(5), 246–249.
28. **Alghadir, Ahmad; Anwer, Shahnawaz; Brismée, Jean-Michel (2015)**. The reliability and minimal detectable change of Timed Up and Go test in individuals with grade 1 – 3 knee osteoarthritis. *BMC Musculoskeletal Disorders*, 16(1), 174–181.
29. **F Salaffi; G Leardini; B Canesi; A Mannoni; A Fioravanti (2003)**. Reliability and validity of the Western Ontario and McMaster Universities (WOMAC) Osteoarthritis Index in Italian patients with osteoarthritis of the knee. , 11(8), 551–560.
30. **Pratibha Prabha et al (2019)**. Efficacy of Russian Current on Pain, Strength of Quadriceps and Function in Subjects with Primary Knee Osteoarthritis: A Randomized Clinical Trial, *International Journal of Health Sciences & Research*, 9(8), 140-148.
31. **John T. Farrar; Andrea B. Troxel; Colin Stott; Paul Duncombe; Mark P. Jensen (2008)**, Validity, reliability, and clinical importance of change in a 0—10 numeric rating scale measure of spasticity: a post hoc analysis of a randomized, double-blind, placebo-controlled trial, 30(5), 0–985.
32. **Akanksha A. Nalawade, Poonam H. Patil, (2020)**, Comparison between Surged Faradic Current and Transcutaneous Electrical Nerve Stimulation (TENS) on Myofascial Trigger Points in Trapezius, *Indian Journal of Physiotherapy and Occupational Therapy*. Vol. 14, No. 1.
33. **S. Epskamp et al (2020)**, Range of motion as an outcome measure for knee osteoarthritis interventions in clinical trials: an integrated review, *Physical Therapy Reviews*, 25:5-6, 462-481.
34. **Norkin C. C., D. J. White (2016)**, Measurement of joint motion: a guide to goniometry, Philadelphia: F.A. Davis Company, Fifth edition, Chapter 1:318-320.
35. **Podsiadlo D, et al (1991)**. *J Am Geriatr Soc*. 1991; 39:142–148.
36. **Shumway-Cook A, et al (2000)**. *Phys Ther.*; 80:896–903.
37. **Bellamy N, Buchanan WW, Goldsmith CH, et al (1988)**: Validation study of WOMAC: a health status instrument for measuring clinically important patient relevant outcomes to antirheumatic drug therapy in patients with osteoarthritis of the hip or knee. *J Rheumatol*, 15:1833-1840.

38. **Michael G. Parkar et al.** Torque Responses in Human Quadriceps to Burst –modulated Alternating Current at 3 Carrier Frequencies. *Journal of Orthopedic and Sport Physical Therapy*, April 2005; 35 (4): 239-245.
39. **Van Baar ME, Assendelft WJ, Dekker J, et al.** Effectiveness of exercise therapy in patients with osteoarthritis of the hip or knee: a systematic review of randomized clinical trials. *Arthritis Rheum.* 1999; 42: 1361–1369.
40. **C. Mang, J. Clair, and D. Collins,** “Neuromuscular electrical stimulation has a global effect on corticospinal excitability for leg muscles and a focused effect for hand muscles.” *Experimental Brain Research*, 209(3): 355-363, 2011.
41. **S.C. Petterson, P. Barrance, T. Buchanan, et al,** “Mechanisms underlying quadriceps weakness in knee osteoarthritis.” *Medicine and Science in Sports and Exercise*, 40(3):422-427, 2008.